

# Ursidae: The Undergraduate Research Journal at the University of Northern Colorado

---

Volume 5

Number 1 2015 Research Excellence Awards Winners  
and Finalists

Article 3

---

January 2015

## Overexpression of the BCL11A Transcription Factor in Basal-like Versus Luminal Breast Cancer Cells

Alexis Cororan

Matthew DeSelm

Follow this and additional works at: <http://digscholarship.unco.edu/urj>

 Part of the [Physical Sciences and Mathematics Commons](#)

---

### Recommended Citation

Cororan, Alexis and DeSelm, Matthew (2015) "Overexpression of the BCL11A Transcription Factor in Basal-like Versus Luminal Breast Cancer Cells," *Ursidae: The Undergraduate Research Journal at the University of Northern Colorado*: Vol. 5 : No. 1 , Article 3.  
Available at: <http://digscholarship.unco.edu/urj/vol5/iss1/3>

This Article is brought to you for free and open access by Scholarship & Creative Works @ Digital UNC. It has been accepted for inclusion in Ursidae: The Undergraduate Research Journal at the University of Northern Colorado by an authorized editor of Scholarship & Creative Works @ Digital UNC. For more information, please contact [Jane.Monson@unco.edu](mailto:Jane.Monson@unco.edu).

## **Overexpression of the BCL11A Transcription Factor in Basal-like Versus Luminal Breast Cancer Cells**

Presenters: Alexis Cororan and Matthew DeSelm

*Faculty Mentor: James Haughian*

**Abstract:** Gene expression profiling identifies discrete subtypes of breast cancers such as luminal, basal-like, and HER2+. Luminal breast cancers respond well to anti-hormonal therapies, and express estrogen (ER) and/or progesterone (PR) receptors. ER, PR, and HER2 are absent in basal-like or triple negative breast cancers, rendering them less responsive to anti-hormonal therapies. Forty percent of luminal breast tumors contain ‘luminobasal’ cells that express the marker cytokeratin 5 (CK5) commonly used to identify basal-like breast cancers. Here we ask- what are these luminobasal cells in luminal tumors? Our models include a pure luminal cell line (E-3) and a pure luminobasal cell line (EWD-8) derived from a parental luminal breast cancer cell line grown in mice. Guided by large microarray datasets comparing women’s primary luminal and basal-like breast cancers, we chose to examine the expression profile of BCL11A between our E-3 and EWD-8 breast cancer cell lines. BCL11A is a transcription factor whose function within breast cancer cells is not fully known; however, microarray profiling indicates that the EWD-8 luminobasal cells express significantly higher levels of BCL11A mRNA relative to luminal cells. Our data confirms this overexpression of BCL11A in the EWD-8 cells, and is associated with the differential expression of accepted luminal and basal-like markers. Future studies will focus on knocking out a specific exon from the BCL11A gene using the Crispr/Cas9 system and assessing the effects in luminobasal cells.