University of Northern Colorado

Scholarship & Creative Works @ Digital UNC

Dissertations

Student Research

12-2019

Psychometric Evaluation of the Casey-Fink Graduate Nurse Experience Survey

Kathryn C. Casey

Follow this and additional works at: <https://digscholarship.unco.edu/dissertations>

Recommended Citation

Casey, Kathryn C., "Psychometric Evaluation of the Casey-Fink Graduate Nurse Experience Survey" (2019). *Dissertations*. 623.

<https://digscholarship.unco.edu/dissertations/623>

This Text is brought to you for free and open access by the Student Research at Scholarship & Creative Works @ Digital UNC. It has been accepted for inclusion in Dissertations by an authorized administrator of Scholarship & Creative Works @ Digital UNC. For more information, please contact Jane.Monson@unco.edu.

UNIVERSITY OF NORTHERN COLORADO

Greeley, Colorado

The Graduate School

PSYCHOMETRIC EVALUATION OF THE CASEY-FINK
GRADUATE NURSE EXPERIENCE SURVEY

A Dissertation Submitted in Partial Fulfillment
of the Requirements for the Degree of
Doctor of Philosophy

Kathryn C. Casey

College of Natural and Health Sciences
School of Nursing
Nursing Education

December 2019

This Dissertation by: Kathryn C. Casey

Entitled: *Psychometric Evaluation of the Casey-Fink Graduate Nurse Experience Survey*

has been approved as meeting the requirement for the Degree of Doctor of Philosophy in the College of Natural and Health Sciences in the School of Nursing, Program of Nursing Education.

Accepted by the Doctoral Committee

Kathleen Dunemn, Ph.D., APRN, CNM, Research Advisor

Melissa Henry, Ph.D., RN, FNP-C, Committee Member

Regina M. Fink, Ph.D., APRN, AOCN, CHPN, FAAN, Committee Member

Chia-Lin Tsai, Ph.D., Faculty Representative

Date of Dissertation Defense October 29, 2019

Accepted by the Graduate School

Cindy Wesley, Ph.D.
Interim Associate Provost and Dean
Graduate School and International Admissions

ABSTRACT

Casey, Kathryn C. *Psychometric Evaluation of the Casey-Fink Graduate Nurse Experience Survey*. Published Doctor of Philosophy dissertation, University of Northern Colorado, 2019.

Graduate nurses experience stress transitioning from the role of nursing student to practicing professional nurse. Understanding the graduate nurse's role transition experience is important because excessive and unrelieved stress might influence job satisfaction and contribute to high turnover. Measuring and improving the role transition experience for graduate nurses has become important for healthcare organizations due to concerns regarding retention. A valid and reliable instrument was needed to assess the graduate nurse's role transition experience. The Casey-Fink Graduate Nurse Experience Survey (Casey, Fink, Krugman, & Propst, 2004) is the most commonly used instrument to measure the stressors, fears, and challenges experienced by graduate nurses during their first year of professional practice.

A secondary data analysis study was conducted to evaluate the psychometric properties of the Casey-Fink Graduate Nurse Experience Survey (Casey et al., 2004). The study sample comprised 71,919 newly graduated nurses who completed the Casey-Fink Graduate Nurse Experience Survey six-months post-entry into the Vizient/American Association of Colleges of Nursing (Vizient/AACN, 2018) one-year nurse residency program. Exploratory factor analysis using principle axis factoring with Promax© rotation was used to determine the underlying factor structure of the set of variables. A five-factor solution yielded a clear pattern of item loadings. This solution accounted for

49.5% of the total variance between items. The five factors were labeled Job Satisfaction, Support, Role Confidence, Organize/Prioritize Care, and Professional Socialization. Confirmatory factor analysis confirmed the five-factor model and cross-validation indicated no revisions were needed to the model. Estimates of internal consistency reliability for the five factors ranged from .73 to .94. This study provided new information to the body of nursing literature on the psychometric testing of the Casey-Fink Graduate Nurse Experience Survey.

ACKNOWLEDGEMENTS

I would like to thank my committee members for their support, encouragement, and assistance during the course of my doctoral research. A special thank you is given to Dr. Chia-Lin Tsai for her expert guidance through the complex factor analysis process. To my dear friend Dr. Regina Fink, I appreciate the many years of friendship we have had. I thank you for understanding me, for sharing your knowledge and experience, and for your confidence in my abilities.

Thank you to my family, friends, and colleagues who have supported me through every phase of my doctoral studies and my professional career. I have felt your steadfast love and support. To my husband Paul, my daughters Claire and Annie, and my parents Arvid and Carol, you have been the wind beneath my wings lifting me along this lifelong journey of learning, and curiosity.

TABLE OF CONTENTS

CHAPTER I. INTRODUCTION TO THE STUDY	1
Background to the Study	2
Significance of the Study	6
The Problem Statement	6
Purpose of the Study	7
Research Question	8
Definition of Terms	8
Summary	8
CHAPTER II. LITERATURE REVIEW	9
Search Strategy	9
Theoretical Frameworks	10
Discussion of Relevant Findings	14
Factor Analysis of the Casey-Fink Survey	19
Summary	22
CHAPTER III. METHODS AND PROCEDURES	24
Research Design	24
Sample Population	24
Sample Size	25
Inclusion and Exclusion Criteria	26
Ethical Considerations	26
Data Collection and Handling Procedures	26
Instrument	28
Data Analysis	29
Limitations of the Study	32
Summary	33
CHAPTER IV. RESULTS	34
Description of the Sample	34
Analysis of Variables	39
Exploratory Factor Analysis	42
Confirmatory Factor Analysis	52

Reliability.....	66
Summary	67
CHAPTER V. DISCUSSION AND CONCLUSION	69
Discussion	69
Major Findings.....	70
Strengths and Limitations	73
Recommendations for Research	75
Conclusion	76
REFERENCES	78
APPENDIX A. INSTITUTIONAL REVIEW BOARD APPROVAL.....	87
APPENDIX B. NON-DISCLOSURE AND CONFIDENTIALITY AGREEMENT	89
APPENDIX C. CASEY-FINK GRADUATE NURSE EXPERIENCE SURVEY	92

LIST OF TABLES

1.	Summary of Casey-Fink Survey Factor Analysis.....	22
2.	Participants by Cohort Year.....	35
3.	Demographic Characteristics of the Six-Month Samples.....	36
4.	Clinical Unit of Employment for the Six-Month Samples	38
5.	Item Means for Section II for Each Sample in the Study	40
6.	Causes of Stress	41
7.	Item Means for Section III for Each Sample in the Study	42
8.	Exploratory Factor Analysis Five-Factor Loadings for Items in Section II	49
9.	Summary of Factor Analysis Models for Section II	50
10.	Exploratory Factor Analysis Loadings for Items in Section III.....	51
11.	Confirmatory Factor Analysis-1 Model Output and Actions Taken	56
12.	Results of Kaiser-Meyer-Olkin Index of Sampling Adequacy and Bartlett's Test of Sphericity.....	58
13.	Communality Scores	59
14.	Pattern Matrix for the Final Five-Factor Solution	62
15.	Confirmatory Factor Analysis-1 Results Using 24 items and Six-Month Sample	65
16.	Reliability Estimates for the Final Five Subscales	67

LIST OF FIGURES

1.	Confirmatory factor analysis-1 five-factor model for 26 items in section II...	53
2.	Confirmatory factor analysis-1 model for items in section III	54
3.	Scree plot for final exploratory factor analysis.....	60
4.	Model used for confirmatory factor analysis-1 analysis.....	63
5.	Confirmatory factor analysis-2 final five-factor structure	66

CHAPTER I

INTRODUCTION TO THE STUDY

Today's healthcare environment is increasingly complex, requiring graduate nurses to acquire new knowledge, abilities, and skills to practice safely and competently. It is very difficult for colleges and schools of nursing to prepare nursing students with the capacity needed to practice in acute care settings immediately after graduation (Goode, Lynn, McElroy, Bednash, & Murray, 2013). Healthcare organizations across the globe are facing significant nurse turnover and are looking for strategies to retain a pipeline of graduate nurses who are eager to join the professional nursing workforce. Kovner and Djukic (2009) reported 26% of graduate nurses leave their first job within two years of starting. In 2014, Kovner, Brewer, Fatehi, and Jun reported 17.5% of newly licensed nurses left their first job within the first year of practice. In an attempt to retain these nurses, healthcare institutions are following the Institute of Medicine (2011) report, *The Future of Nursing: Leading Change, Advancing Health*, which recommended the implementation of nurse residency programs to facilitate transition to practice and to reduce nurse turnover. These programs have been developed and embraced as an innovative intervention to provide support for newly graduated nurses as they transition into practice, to increase clinical competency in the professional role, to improve confidence and job satisfaction, and to improve graduate nurse retention (Goode et al., 2013; Goode, Ponte, & Havens, 2016; Krugman et al., 2006).

There is current interest in evaluating the effectiveness of transition into practice programs. The American Nurses Credentialing Center's (ANCC, 2018) Practice Transitions Accreditation Program was established to ensure these programs conformed to evidence-based standards. The Commission on Collegiate Nursing Education (CCNE, 2015) created an accreditation process for postbaccalaureate nurse residency programs that was developed in partnership with academic nursing programs. These accreditation standards all required current, reliable, and valid evaluation methods to demonstrate program quality and to measure outcomes (Stephenson & Cosme, 2018). Using a valid measurement instrument to evaluate key concepts related to graduate nurse role transition, one that has been rigorously tested and reflects the current nursing practice environment, could provide evidence for nurse residency program design and improve effectiveness in supporting transition to the profession.

Background to the Study

Transitioning from the role of nursing student to professional nurse is a period of stress, role adjustment, and reality shock (Casey, Fink, Krugman, & Propst, 2004). The new nurse is often confronted with the realization of being unprepared for their new role and responsibilities, which often results in job stress, job dissatisfaction, and thoughts of leaving nursing (Brewer, Kovner, Greene, Tukov-Shuser, & Djukic, 2012). Casey et al. (2004) found graduate nurses reported a lack of confidence in skill performance, deficits in critical thinking and clinical knowledge, struggles with peer and preceptor relationships, and dependence on others. In addition, the desire to be independent practitioners, frustrations with the work environment, lack of organizational skills, and communication with physicians were difficulties experienced during role transition.

Graduate nurses are becoming a significant portion of hospital new hires and staffing strategies as nursing workforce shortages become a reality; yet, graduate nurse turnover rates have been reported between 30% and 60% (Bowles & Candela, 2005; Harrison & Ledbetter, 2014).

Measurement of graduate nurse role comfort, confidence, job satisfaction, and skill acquisition during this transition into professional practice is essential and must be assessed accurately and consistently using a carefully constructed and validated scale. The measurement tool most frequently used in published studies (e.g., Anderson, Hair, & Todero, 2012; Letourneau & Fater, 2015) to measure graduate nurse self-reported perceptions of the transition to practice experience is the Casey-Fink Graduate Nurse Experience Survey (Casey et al., 2004). Measuring the role transition experience of graduate nurses is needed to assist in improving the curriculum and effectiveness of transition to practice programs and in the retention of graduate nurses new to the practice setting.

This cross-sectional study used a secondary data analysis design to evaluate the psychometric properties of the Casey-Fink Graduate Nurse Experience Survey (Casey-Fink Survey; Casey et al., 2004). Factor analysis is a useful approach to assessing construct validity when an investigator has designed a measure to assess various dimensions or sub-components of a phenomenon of interest and wishes to empirically justify these dimensions or factors (Soeken, 2010). The data set for this study was collected using the Casey-Fink Survey from a convenience sample of graduate nurses who participated in the Vizient/American Association of Colleges of Nursing (Vizient, 2018) one-year post-baccalaureate nurse residency program during a 10-year span of time

(January 2008 to December 2018). The sample included newly graduated nurses with less than one year of clinical experience. The Casey-Fink Survey is a self-report instrument designed to measure new nurses' perceptions of role transition issues experienced at entry into practice and through the first 12 months of professional practice. An exploratory factor analysis, confirmatory factor analysis, and reliability testing were conducted to establish current psychometric properties of the Casey-Fink Survey and further refine the factors that influenced role transition.

This researcher and Regina Fink (Ph.D., RN, AOCN, FAAN) co-developed the initial Casey-Fink Survey (Casey et al., 2004) instrument in the spring of 1999. Informal feedback from former students and observed turnover patterns of graduate nurses motivated the authors to design this measurement survey. Conceptualization of the items for the survey resulted from a comprehensive literature review and clinical practice experience of the authors that assisted in identifying constructs to study. Relevant theoretical concepts from Patricia Benner's (1984) *Novice to Expert Theory of Skill Acquisition* and Marlene Kramer's (1974) *Reality Shock* theory were used as the frameworks to guide item development for this evaluative measurement tool. Survey items were selected and developed as indicators of the underlying construct: role transition. Various response formats were used in the development of the survey. Responses to the items in section II of the survey were formatted using a 4-point Likert scale response, omitting a neutral response option. Items in section III were formatted using a 5-point Likert scale response.

The instrument was reviewed for face and content validity using an expert panel of nursing practice leaders and academic educators employed at the University of

Colorado College of Nursing and at a large academic medical center prior to being pilot tested on 12 graduate nurses employed at an urban academic medical center (Casey et al., 2004). After the initial questions were pretested and insights from potential respondents were gathered, a longitudinal study was initiated to describe graduate nurses' experiences at specific time periods as they transitioned into the professional nurse role. The Casey-Fink Survey (Casey et al., 2004) was administered to a convenience sample of 270 graduate nurses working in six acute care urban hospitals located in the Denver-Metro area. Internal consistency reliability was established with Cronbach's alpha of .78 on the 24 items in section II with Likert scale responses of 1= *Strongly disagree* to 4= *Strongly agree*). Questions in section II related to graduate nurse perceptions of their levels of comfort, confidence, and support while functioning as a professional nurse. Frequencies for responses to skills uncomfortable performing independently, satisfaction with aspects of their job, difficulties with role transition, and demographics of the respondents were reported. Based on responses from the initial sample of graduate nurses, the survey was revised in 2002 to include additional questions about the work environment. The survey was revised a second time in 2006 following results of a qualitative analysis that permitted converting the open-ended items in section IV to a multiple-choice format (Fink, Krugman, Casey, & Goode, 2008). After these revisions, internal consistency reliability remained unchanged. In 2004, a factor analysis was completed on section II of the survey. In the analysis, a 5-factor solution was found, accounting for 46% of the variation in total scores. The factors were labeled Support, Patient Safety (Organizing/Prioritizing), Stress, Communication/Leadership, and Professional

Satisfaction. Reliability estimates for the factors ranged from .71-.90 (M. Lynn, personal communication, July 10, 2004).

Significance of the Study

Graduate nurse transition has been the focus of attention for nurse leaders in both academic and practice institutions for over 40 years since the publication of the seminal work: *Reality Shock* (Kramer, 1974). Understanding current difficulties new nurses face during the transition period might uncover unique factors that influence retention and job satisfaction. Graduate nurses are becoming the employment pipeline for acute care hospitals. These newly graduated nurses are entering the workforce and finding they have neither the practice expertise nor the confidence to navigate what has become a highly dynamic and intense clinical environment burdened by escalating levels of patient acuity and nursing workload (Duchscher, 2008). A major risk with role transition is high stress and turnover rates experienced by graduate nurses during their first year of professional practice. A valid and reliable survey instrument is required to measure role transition constructs identified in theoretical frameworks of transition. Using a standard measurement tool allows healthcare organizations to compare their results with other organizations using the same population of interest. This study evaluated the psychometric properties of the Casey-Fink Survey and tested the accuracy of the theoretical concepts proposed in the theories selected for use in this study.

The Problem Statement

The Casey-Fink Survey has been widely used as a valid tool to assess the graduate nurse's transition into practice experience and as a program evaluation measurement tool for nurse residency/orientation programs (Fink et al., 2008). As the Casey-Fink Survey

was constructed over 20 years ago, a re-evaluation was needed given the changes in educational programs and clinical practice over this extended time period. In addition, the length of the survey might contribute to survey fatigue when administered over three time periods. Thus, a shorter survey might be appropriate and should be considered. The factor structure of the scale in section II was last examined in 2004 and a comprehensive psychometric evaluation of the survey instrument was overdue. A construct validation study using a large sample of subjects was the best means of demonstrating the tool's usefulness in measuring constructs of role transition and the experiences of a graduate nurse's role transition into the practice setting. Ideally, scale developers should take steps to gather new data about the worth of their instrument in a validation study (Polit & Beck, 2012).

Purpose of the Study

The purpose of this descriptive study was to evaluate the psychometric properties of the Casey-Fink Survey, a self-report instrument, used to measure perceptions of role transition factors newly graduate nurses experience during the first 12 months of professional practice. Testing of the instrument included examination of its structure, reliability, and construct validity. A secondary analysis of data collected using the Casey-Fink Survey with a large sample size helped assess the validity and reliability of the measure, the trustworthiness of the findings, and contributed to generalizable knowledge on the concepts being measured by the Casey-Fink Survey.

Research Question

The following research question guided this study:

- Q1 What role transition constructs are measured by the Casey-Fink Survey in newly graduated nurses who participated in the 1-year Vizient/AACNTM Nurse Residency Program?

The specific aims of this study were to

1. Explore the factor structure of the Casey-Fink Survey,
2. Test the structural validity of the Casey-Fink Survey, and
3. Assess the reliability of the Casey-Fink Survey.

Definition of Terms

Graduate nurse. One who has graduated from a basic nursing education program, passed the licensing exam, and has no professional nursing experience. Graduate nurses remain in this role for a period of 6 to 12 months as defined by the hiring institution.

Role transition. The experiences of moving from the student nurse role into a professional nurse role. Role transition was measured using the Casey-Fink Survey.

Summary

This chapter described the study including the background, significance, and problem statement. Definitions of key terms used in the study were included in this chapter. A psychometric evaluation to estimate the reliability and examine structural validity of the Casey-Fink Survey would be a significant contribution to the body of nursing knowledge on measurement instruments specific to graduate nurse role transition. In addition, this study was needed to inform revisions to the Casey-Fink Survey.

CHAPTER II

LITERATURE REVIEW

Chapter II outlines the search strategy used for the literature review, the theoretical frameworks guiding measurement of constructs in this study, relevant findings reported related to measurement of the constructs associated with graduate nurse role transition using the Casey-Fink Survey, and findings related to evaluating the psychometric properties of the Casey-Fink Survey are presented. In the conclusion, a summary highlighting the research gap is provided.

Search Strategy

An initial search of relevant literature published from 2000 to 2018 was completed using the Cumulative Index to Nursing and Allied Health Literature (CINAHL) and PubMed databases. Key words included “Casey-Fink Graduate Nurse Experience Survey,” “new graduate nurse,” “transition to practice,” “role transition,” “transition”, “newly licensed nurse”, “graduate nurse”, “instrument development” and “instrument validation.” Summon, CINAHL, and PubMed were further searched using the following terms: “Casey-Fink Graduate Nurse Experience Survey” AND “factor analysis.” Two published dissertations reporting a factor analysis on the Casey-Fink Survey were found. All titles and abstracts were reviewed for the following inclusion criteria: English language, full text, dissertations, and peer-reviewed journal articles. Studies addressing graduate nurse role transition using the Casey-Fink Survey as one of

the measurement instruments were included in this literature search and are reviewed here.

Theoretical Frameworks

Four frameworks provided the theoretical basis for this research study. Meleis' (2010) *Theory of Nursing Transitions* proposed that assisting people to manage life transitions is a key function of nursing (Schumacher & Meleis, 1994). A transition is defined as a "passage or movement from one fairly stable state, condition, or place to another fairly stable state, and is a process triggered by change" (Meleis, 2010, p. 11). According to this framework, role transition for graduate nurses into professional practice is considered a situational transition. The transitions model includes four defining concepts: the nature of transitions, transition conditions, patterns of response, and nursing therapeutics (Meleis, Sawyer, Im, Hilfinger, & Schumacher, 2000). Transitions theory offers a framework for understanding the graduate nurse's experience of transition, for uncovering the conditions that facilitated and hindered a successful transition experience, for describing their responses during the transition, and in identifying strategies or interventions that contributed to successful outcomes of transition. Three conditions influence role transition: personal conditions, community conditions, and societal conditions. Transitions theory identifies nurses as having a key role in enhancing a sense of well-being in their patients. Transitions often require a person to incorporate new knowledge, to alter behavior, and, therefore, to change the definition of self in a new social context (Meleis et al., 2000). The challenge for nurses and others involved in supporting those undergoing a transition is to understand the transition process and develop interventions that are effective in helping them regain stability and a sense of

well-being (Schumacher & Meleis, 1994). Transition involves new knowledge, change in behaviors, and requires support to regain a sense of well-being (Meleis, 2010).

According to this model, a transition process is triggered by the event of starting a new job as a graduate nurse, which prompts a need for support and assistance in learning effective strategies to cope with this change. Meleis found outcomes of successful transition were developing confidence, coping, feeling connected, and skill mastery.

Benner's (1984) novice to expert theory of skill acquisition articulated stages of clinical competence in nursing practice. Benner's theory, as applied to the nursing profession, was adapted from the Dreyfus model of skill acquisition (Benner, 1984). According to this model, graduate nurses who enter the nursing profession are usually at the advanced beginner stage. They develop knowledge and acquire skills in an incremental manner while moving along the continuum from novice to advanced beginner. The advanced beginner nurse demonstrates marginally acceptable nursing performance and gains experience in specific and actual situations (Benner, 1984). Advanced beginners benefit from having a preceptor or experienced nurse to provide guidance, mentoring, and to explain recurring and meaningful components of situations (Benner, 1984). The transition from student to new graduate nurse marks the beginning of the journey from novice to advanced beginner (Marshburn, 2007).

Kramer (1974) outlined a theory of newly graduated nurse transition that identified specific stages and characteristics of role transition and proposed strategies for providing support. The term "reality shock" was used to describe the phenomenon and the specific shock-like reactions of new nurses when they found themselves in a work situation for which they thought they were prepared and then suddenly found they were

not (Kramer, 1974). This shock-like reaction followed when the newly graduated nurse perceived that many ideals and values taught in the educational program were not operationalized and went unrewarded in the work setting (Kramer, 1974). Kramer theorized that those new to the nursing profession went through four stages of transition: (a) the honeymoon phase—a period of excitement to be joining the profession; (b) the shock phase—a period when negative feelings toward the profession surfaced, when the new nurse was most vulnerable, and at risk to quit or leave their unit; (c) the recovery phase—when the new nurse was able to view the job realities with a more open perspective, and lastly (d) the resolution phase—usually occurred around one year when the new nurse could see the professional role in perspective and could fully contribute to the profession (Kramer, 1974). This role transformation process meant the new nurse separate themselves from the expectations held for them in school and, concomitantly, took on new expectations that arose in the context of their new jobs (Kramer, 1974, p. 192). Having self confidence in one's ability to learn and use one's resources is a primary requisite for making a smooth transition from the school environment to the work setting (Kramer, 1974). Kramer was the first to call for structural changes to better support graduate nurses during the role transformation process and outlined an effective program of socializing new nurses into the attitudes, expectations, and behaviors of the professional work setting so the new nurse learned to perform his/her role effectively.

Duchscher's (2008) theory of transition shock built on Kramer's (1974) reality shock theory. Using a grounded theory process, Duchscher described graduate nurse role transition as progressing through three main stages: doing, being, and knowing. Duchscher learned through her qualitative work that in the doing phase, graduate nurses

had idealistic expectations and anticipations that were far from reality. The nurses blamed this disparity on a lack of educational preparation (Duchscher, 2008). After a few months in practice, they reported feeling more comfortable, had accepted their limitations, could ask questions of their colleagues, and were then mentally ready to move to the next phase of learning (Duchscher, 2008). In the second phase (being), graduate nurses had an increase in knowledge level, skill competency, and critical thinking while they began settling into their new role and responsibilities. During this stage, new nurses gained trust and confidence in their own capabilities and began to apply practical meaning to their theoretical knowledge (Duchscher, 2008). In the final stage—knowing, the new nurse sees a shift in personal and professional socialization, views the influences of stress on themselves, and moves from a position of insecurities and abilities to frustrations with the system and being at the bottom of the pecking order (Duchscher, 2008). Duchscher described this as transition crisis. The theoretical construct of transition shock demonstrated the newly graduated nurse, engaging in a professional practice role for the first time, was confronted with a broad range of physical, emotional, developmental, intellectual, and sociocultural changes that were factors due to the experience of transition (Duchscher & Windey, 2018). The stages of professional role transition reflect a non-linear process that moves the new practitioner through developmental and professional, intellectual and emotive, skill and role-relationship changes, and contains within it experiences, meanings, and expectations (Duchscher, 2008). The healthiest transition experiences for graduate nurses would be facilitated when they had personal and professional lives characterized by stable and supportive relationships, were afforded roles and responsibilities commensurate with

their stage of transition, received consistent workplace support and constructive feedback, were familiar with care delivery and skill performance, were provided opportunities to be supported by experienced nurses about increasingly complex clinical decisions and judgements, and were supported in their workplace environments (Duchscher, 2008).

Based on Benner's (1984) and Duchscher's (2008) theories, graduate nurses experience an increase in knowledge, stress, and confidence during their initial six months of practice and an increase in communication and professional satisfaction at the end of the first year of practice. Kramer's (1974) four stages of reality shock provided a basis to explore the graduate nurse experience during the first year of transition to practice. These four time points were used by Casey et al. (2004) to survey graduate nurses' perceived levels of confidence and comfort in the first year of professional practice. Confidence and comfort in the professional nurse role, supportive relationships, presence of stressors, stages of skill and knowledge acquisition, and changes in self are common attributes and characteristics of role transition identified in all four theoretical frameworks. These theoretical frameworks offered empirical indicators of the selected concepts of interest and provided a foundation for studying the constructs being measured in the Casey-Fink Survey (Casey et al., 2004).

Discussion of Relevant Findings

Twenty descriptive studies from the literature were reviewed where the Casey-Fink Survey was used to collect data. Nineteen studies were based in the United States and one was in British Columbia. All studies were published between 2009 and 2018, indicating an increased interest in measuring graduate nurse role transition since the Institute of Medicine's (2011) report: *The Future of Nursing: Leading Change*,

Advancing Health. Fourteen studies used a pretest/posttest design using repeated measures analysis and did not have a comparison group (Cline, La Frentz, Fellman, Summers, & Brassil, 2017; Dubois & Zedreck Gonzales, 2018; Friday, Zoller, Hollerbach, Jones, & Knofczynski, 2015; Goode, Lynn, Krsek, & Bednash, 2009; Goode et al., 2013; Klingbeil et al., 2016; Medas et al., 2015; Meyer & Shatto, 2017; Olson-Sitki, Wendler, & Forbes, 2012; Slate, Stavarski, Romig, & Thacker, 2018; Thomson, 2011; Tyo et al., 2018; Wenger, 2015; Wilson, Weathers, & Forneris, 2018). No studies used a randomized controlled trial design. Two studies used a pre-test design (Cylke, 2012; Marshburn, Engelke, & Swanson, 2009). Four studies used a posttest design (Harrison & Ledbetter, 2014; Hillman & Foster, 2011; Johnson, Salisbury, Johannsson, & Barajas, 2013; Rush, Adamack, Janke, Gordon, & Ghement, 2012). Sample sizes in these studies ranged from 11 to 1,638. All studies used convenience samples of graduate nurses who were employed in academic medical centers.

Five studies used section II of the Casey-Fink Survey (Fink et al., 2008) exclusively to evaluate their nurse residency program outcomes and to inform changes in the program curriculum. The exclusive use of section II, which measured the concepts of support, patient safety, stress, professional satisfaction, and communication/leadership, was the most common survey modification reported in studies from this review of the literature. The Thomson (2011) study used sections I, II and V of the Casey-Fink Survey to compare subscale scores of associate degree and baccalaureate degree nurses within the same organization who participated in the same program curriculum. Results reported differences in support and professional satisfaction between the two groups (Thomson, 2011). Two studies used questions from section II of the survey to evaluate a

graduate nurse resilience training program (Dubois & Zedreck Gonzales, 2018) and to compare the relationship between transition to practice experiences and resiliency (Meyer & Shatto, 2017). Wenger (2015) used skills in section I to demonstrate comfort performing intravenous (IV) skills independently. This study demonstrated improved confidence in performing IV skills from baseline to six months of practice following an IV insertion class provided to graduate nurses in the hiring institution.

Two studies reported retrospective analysis of 10 years of data collected using the Casey-Fink Survey. Cline et al. (2017) conducted a retrospective analysis of 10 years of data collected using the Casey-Fink Survey from graduate nurses who participated in an internally developed residency program. A total of 1,638 graduate nurses completed surveys at three time periods and were included for analysis. Pre-participation and post-participation scores were compared using a two-sample *t*-test. Results indicated statistically significant changes in scores for all factors except stress.

Communication/leadership and patient safety (organizing/prioritizing) scores demonstrated the most favorable improvement with an increase in mean score from 2.88 to 3.24 ($p < .001$), indicating increased confidence and comfort in communicating with interprofessional colleagues, patients, and families and in delegating skills (Cline et al., 2017). The patient safety domain increased from 2.77 to 3.16 ($p < .001$), indicating improved confidence in organizing and prioritizing patient care and comfort with safely completing the components of the patient care assignment. Scores in the support and professional satisfaction domains showed a moderate but significant decline during the course of the residency program. Mean scores in the support domain decreased from 3.36 to 3.29 ($p = .002$) and mean scores in the professional satisfaction domain decreased

from 3.53 to 3.41 ($p < .001$; Cline et al., 2017). Stress responses in this study ranged from 0.00 to 1.00, reflecting a low level of stress among residency participants. Changes in the mean scores from 0.15 to 0.13 suggested a small decrease in stress that approached significance ($p = .05$; Cline et al., 2017). Graduate nurse retention over the 10 years reflected a high retention rate at $>90\%$ at one year. Cline et al. (2017) stated variability in the survey questions asked and a changing residency program curriculum that evolved over time to meet the emerging needs of the institution and newly hired nurses were cited as limitations to this study. Strengths of this study suggested internally developed residency programs might be equally effective as prepackaged programming in supporting graduate nurse safety, confidence, and retention of these newly licensed nurses (Cline et al., 2017).

Goode et al. (2013) examined outcomes from 10 years of a post-baccalaureate nurse residency program. The researchers utilized the Casey-Fink Survey (Casey et al., 2004) to collect and analyze data from graduate nurses participating in the University Health System Consortium/American Association of Colleges of Nursing Nurse Residency Program from 2002 through 2012. This study examined perceptions of 1,016 graduate nurses at the beginning, at six months, and end of the one-year program. Repeated measures analysis of variance (ANOVA) found significant increases from program start to program completion in graduate nurses' perceptions of their overall confidence, support, ability to organize and prioritize their work, and communicate and provide clinical leadership. Professional job satisfaction showed a significant decline from program start to the six-month point and then stabilized. Job stress increased from beginning to the end of the program. The reason for the change in graduate nurses'

scores of job stress was not discussed in their article. Limitations of this study included lack of a control group, difficulty to maintain a high response rate in repeated measures ANOVA, and not being able to control for individual site related extraneous variables (Goode et al., 2013).

Reliability testing is a measure of the amount of random error in a measurement technique (Grove, Gray, & Burns, 2015). Cronbach's alpha coefficient is the most commonly used measure of reliability scales with multiple items and estimates of reliability are specific to the sample being tested (Grove et al., 2015). Researchers need to perform reliability testing on each instrument used in a study to ensure it is reliable for that study (Grove et al., 2015). Only three of the studies reported reliability testing (Cylke, 2012; Marshburn et al., 2009; Olson-Sitki et al., 2012).

Slate et al. (2018) reported outcomes of a nurse residency program using the Casey-Fink Survey, which indicated improvements in the onboarding experience for new graduate nurses in their longitudinal study. A convenience sample of 208 graduate nurses completed the survey at two different time points. Analysis of variance was used to compare the number of primary preceptors, which resulted in a decrease in the number of preceptors used to onboard graduate nurses and standardization of the program length. Findings using the Casey-Fink Survey provided insights into the disappointment participants felt related to multiple preceptors with varying effectiveness (Slate et al., 2018). Data from this study informed changes in the onboarding process for new graduates including the numbers of preceptors, program length standardization, and improvements of emergency clinical response education (Slate et al., 2018). Reliability estimates were not completed in this study.

Factor Analysis of the Casey-Fink Survey

A common method for assessing the construct validity of an instrument is exploratory factor analysis (EFA). This analytic tool helps an investigator determine empirically how many constructs, or latent variables, or factors underlie a set of items (DeVellis, 2017). These underlying factors are defined in mathematical terms so the process is considered data-driven (Soeken, 2010). In EFA, the researcher does not know how many factors or latent variables will be formed and uses specific criteria to determine the factors. Thus, EFA is generally described as a descriptive or exploratory procedure. In confirmatory factor analysis (CFA), the intent is to hypothesize or define the factors directly and then determine how well the defined measurement model fit the observed data (Soeken, 2010). Confirmatory factor analysis is often used in later phases after the underlying structure has been established on prior empirical (EFA) and theoretical grounds (Brown, 2015). Confirmatory factor analysis is a theory-driven approach rather than data-driven (Soeken, 2010).

In 2004, items in section two of the Casey-Fink Survey were subjected to exploratory factor analysis—principal axis factoring with Varimax[®] rotation. Principal axis factoring was selected to decrease the likelihood of overestimating the explained variance and item factor loadings common with principal component's analysis (M. Lynn, personal communication, July 10, 2004). In this analysis, a five-factor solution was found, accounting for 46% of the variation in total scores: support, stress, patient safety (organizing/prioritizing), communication/leadership, and professional satisfaction. Reliability estimates for the factors ranged from .71 to .90 (M. Lynn, personal communication, July 10, 2004). The instrument was originally scored by summing all of

the items in section two, including the stress items, for an internal consistency estimate of ($\alpha = .89$). Confirmatory factor analysis was not conducted.

In the Marshburn (2007) study, 265 new nurses comprised the study sample. Principal components analysis (PCA) was conducted on the 24 items from part two of the Casey-Fink instrument. Principal components analysis of the 24 items revealed the presence of five factors with eigenvalues greater than 1.00 (Marshburn, 2007). Inspection of the scree plot revealed a clear break after the four components and the parallel analysis also suggested four factors be retained for further analysis (Marshburn, 2007). Those four components explained 56.9% of the total variance in the 24 items. Varimax rotation was then performed to aid in the interpretation of the four components. An inspection of the rotated component matrix revealed the fourth component had four items with loadings greater than .50 but two of the items had cross-loadings greater than .47 (Marshburn, 2007). Principal components analysis was re-run with only three components extracted; the resulting initial solution was subjected to another Varimax rotation. After inspecting the items that loaded on each component, the first subscale was labeled Support, the second subscale was labeled Patient Care, and the third subscale was labeled Professional Role (Marshburn, 2007, p. 787-788). The three subscales (Support, Patient Care, and Professional Role) derived from the PCA were used as new variables in this study to examine relationships of new nurses' perceptions of clinical competence and measured performance-based clinical competence (Marshburn, 2007). Confirmatory factor analysis was not conducted in this study.

In the Hallaran study (2017), the study sample consisted of 217 new nurses. Exploratory factor analysis of the Casey-Fink survey resulted in the identification of seven factors:

The first factor had items loading from a different subscale. The second factor had four of the five patient safety items as the highest loading items and supported retaining that subscale. A Cronbach's alpha for the revised subscale of four items was .78. The third factor had three items from the Support subscale, each addressing being able to get help and feedback from nursing peers and staff. As these items related to each other, a new subscale was created, Peer Support, and resulted in a Cronbach's alpha of .78. The fourth component had five of the six items from the Communication/Leadership subscale with high loadings. The Cronbach's alpha for the subscale, without one item, was .64. The fifth factor had three items from the original Support subscale, which addressed support from preceptors and role models. As these relationships were different from those of peers, a new subscale was created, Preceptor Support, which had a Cronbach's alpha of .77. The sixth factor contained two of three Professional Satisfaction subscale items and supported a revised subscale. The Cronbach's alpha of the two-item scale was .84. The seventh factor had two items: one stress item (negatively scored) and one professional satisfaction item. A review of the professional satisfaction item revealed the item focused on feeling supported by family and friends. The stress item was specific to personal stress. Therefore, these two items did relate and a revised stress subscale was formed with the

scores of the stress items recoded. The Cronbach's alpha of the revised scale with two items was very low at .32. (Hallaran, 2017, p. 86)

Confirmatory factor analysis was conducted in this study. The seven factors were reduced to two factors. A major limitation of this study was the small sample size. Table 1 provides a summary of the survey factor analysis.

Table 1

Summary of Casey-Fink Survey Factor Analysis

Author Year	Number of Factors Retained	Factor Names & Number of Items	Sub-Scale Cronbach's alpha	Extraction & Rotation Method
Lynn (2004)	5	Support (9 items)	.90	Principal Axis Factoring
		Patient Safety (5 items)	.79	
		Stress (6 items)	.71	
		Communication/Leadership (6 items)	.75	Varimax Rotation
		Professional Satisfaction (3 items)	.83	
Marshburn (2007)	3	Support (11 items)	.89	Principal Component Analysis Varimax Rotation
		Patient Care (4 items)	.73	
		Professional Role (5 items)	.67	
Hallaran (2017)	5	Patient Safety (4 items)	.78	Principal Component Analysis
		Peer Support (3 items)	.78	
		Communication/Leadership (5 items)	.64	
		Preceptor Support (3 items)	.77	Varimax Rotation
		Professional Satisfaction (2 items)	.84	

Summary

Chapter II provided an analysis of four theories of transition that identified constructs to be measured, examined the literature related to studies that measured factors of graduate nurse role transition using the Casey-Fink Survey, and reviewed past testing and psychometric analysis of the Casey-Fink Survey. The Casey-Fink Survey was originally designed to measure concepts related to graduate nurse role transition; yet, it is

often used and revised by investigators to measure outcomes of a nurse residency program. Conducting a psychometric evaluation study using EFA and CFA with a large sample size collected from graduate nurses using the entire Casey-Fink Survey provided an opportunity to test and confirm the factor structure of the survey and to demonstrate the trustworthiness of the findings. A lack of published EFA and CFA results for the Casey-Fink Survey further supported the need for additional testing of the instrument.

CHAPTER III

METHODS AND PROCEDURES

Chapter III provides an overview of the research design, methods, and data analysis procedures. This chapter also includes a description of the study population and sample, sample size, inclusion and exclusion criteria, presentation of ethical considerations and protection of human subjects, data collection and management procedures, description of the measurement instrument, analysis of the data, and limitations of the study.

Research Design

A descriptive cross-sectional study design was used to evaluate the psychometric properties of the Casey-Fink Survey. A secondary analysis of data previously collected from newly graduated nurses who participated in the Vizient/AACN (Vizient, 2018) 12-month Nurse Residency Program from 2008 to 2018 was used to establish construct validity in this study. A secondary analysis was an analysis of information or data previously collected by another researcher or organization (Grove et al., 2015).

Sample Population

This study used an existing database of responses gathered from graduate nurses who participated in the standardized 12-month nurse residency program developed by Vizient/AACN (Vizient, 2018). The sample was obtained using nonprobability, convenience sampling of 10 years of data collected with the Casey-Fink Survey from participants in the Vizient/AACN 12-month Nurse Residency Program. Convenience

sampling is a nonprobability sampling technique in which subjects are included in the study based on their accessibility and proximity (Gray, Grove, & Sutherland, 2017). Study participants included newly graduated nurses who were working in over 450 hospitals in the United States (Vizient, 2018). The Vizient/AACN 12-month Nurse Residency Program offers a standardized monthly transition to practice curriculum including monthly classroom sessions that focus on leadership, patient outcomes, and professional development while providing peer and leadership support (Vizient, 2018). A graduate nurse is defined as one who has graduated from a basic nursing education program, passed the licensing exam, and has no professional nursing experience.

Sample Size

The number of subjects needed to undertake a factor analysis of an instrument depends on the number of items initially included (Pett, Lackey, & Sullivan, 2003). Pett et al. (2003) suggested at least 10 to 15 subjects per initial item, preferably aiming for a sample size of 500 to 1,000 subjects. An estimated sample size of 80,000 was calculated based on obtaining 10 years of data collected and the 2018 Vizient report that on average, 18,000 graduate nurses per year participate in their 12-month program. Participants voluntarily completed the Casey-Fink Survey at beginning of the program, at six months, and at the end of the program. The six-month time period for data analysis was chosen to test the factor structure as this was usually when the graduate nurse had developed an increase in his/her knowledge and skill competency, was settling into his/her new role responsibilities, and had time to adjust to work environment expectations.

Inclusion and Exclusion Criteria

Inclusion criteria were that study participants must be 20 years of age, had less than or equal to six months of professional nursing experience, had graduated from an accredited baccalaureate or associate degree, diploma, or master's degree registered nursing (RN) education program, and participated in the Vizient/AACN (Vizient, 2018) 12-month nurse residency program. The exclusion criterion was those newly licensed nurses who had greater than six months of RN experience. Respondents who did not complete the survey at the six-month time period were excluded from the study.

Ethical Considerations

Compliance with all guidelines and ethical principles for research with human subjects was adhered to as outlined by the Collaborative Institutional Training Initiative (2017) program. There were no foreseeable significant social or psychological risks to the graduate nurses from participation in the study. Risks inherent in this study were no greater than those normally encountered during regular classroom participation. Consent to participate in research was obtained after the graduate nurse logged into a password protected Vizient database at the study site. Potential benefits to the participants included having the satisfaction of providing valuable feedback for designing strategies to support the transition of graduate nurses, develop education needed to be successful in their first job, and to promote the retention of graduate nurses.

Data Collection and Handling Procedures

After Institutional Review Board approval was granted from the University of Northern Colorado (see Appendix A), de-identified participant survey responses (raw data) were sent to the researcher from the Vizient/AACN's Senior Program Manager at

the Vizient corporate office. A data use agreement (see Appendix B) was signed for the release of the data to this researcher. Data were received from Vizient/AACN (Vizient, 2018) through a secure electronic lock box and consisted of a Microsoft 2010 Excel file. The data were imported into the statistical software installed on the researcher's personal password protected computer and was accessible only by the researcher. The researcher's personal computer was protected with the latest version of Norton's Security Suite that was last updated on February 5, 2019 and included a firewall, virus protection, and tamper protection.

The Excel file contained 73 columns of data that included a unique organization identification number, survey period, cohort year, unit, age, gender, ethnicity, degree received, previous non-nursing degree, and participant responses to questions on the Casey-Fink Survey. The original Excel file was retained in a folder on the researcher's computer, filtered to include only the six-month survey period data, imported into Statistical Package for the Social Sciences (SPSS) software version 26, and prepared for data cleaning and analysis in SPSS. A unique identification number was given by the Vizient Senior Program Manager to each organization that participated in the Vizient/AACN 12-month nurse residency program. The key for the unique organization number was not shared with the researcher, thus ensuring maximum confidentiality and anonymity of all participants and study sites. No information contained within the Excel files could be traced back to a study participant or study site.

All data entered into SPSS were examined for accuracy of coding and completeness during the cleaning process to minimize errors in data entry. There were 71,920 rows of data representing 71,920 participant responses in the six-month survey

time period. One participant did not answer any of the survey items and was excluded from the data analysis. An individual identification number was added as a variable in the data set. The age of participant responses for inclusion in the data analysis was 20 to 65 years of age; there were 5,633 missing responses for age. The range of responses was 0 to 223. Age responses entered as 0-19 ($N = 77$) were deleted. Age responses entered as 99-223 ($N = 52$) were deleted. Gender responses were recoded as numeric and retained as female and male. All eight ethnicity categories were re-coded as numeric and reserved for analysis. Nursing degree categories were condensed to five groupings and re-coded as numeric. Previous non-nursing degree was re-coded as numeric and all five categories were retained. All remaining questions in the data set were recoded as numeric. The SPSS data file of 71,919 responses was then split into three samples divided randomly from the entire 6-month data sample to represent the same population. The EFA sample was 33% ($N = 23,489$) of the total sample and was used for the exploratory factor analysis. The second sample labeled CFA-1 was 33% ($N = 24,378$) of the total sample and was used for the confirmatory factor analysis. The third sample labeled CFA-2 was 34% ($N = 24,052$) of the total sample and was used to cross-validate the findings found in the CFA-1 analysis.

Instrument

The Casey-Fink Graduate Nurse Experience Survey (see Appendix C) was selected as the measurement instrument to be evaluated in this study. This instrument was published in 2004 and revised in 2006. The survey includes five sections with a total of 55 questions. Section I includes one question where respondents are asked to select the top three skills or procedures they are most uncomfortable performing independently

using a drop down list of 21 skills. They could also select “I am independent in all skills.” Section II contains 25 items. A response option for each of the first 24 items consists of a 4-point Likert scale asking respondents to indicate their response from the following four choices: *Strongly Disagree*, *Disagree*, *Agree*, and *Strongly Agree*. An additional question (#25) is where the respondent answers “yes” or “no” to a series of six stressors. Section III consists of nine questions designed to rate satisfaction with various aspects of the job. Response options are on a 5-point Likert scale indicating *Very Dissatisfied*, *Moderately Dissatisfied*, *Neither Satisfied nor Dissatisfied*, *Moderately Satisfied*, and *Very Satisfied*. Section IV consists of four multiple choice questions and one open-ended question. The four questions ask participants to circle the responses that apply to their transition experience: (a) “What difficulties, if any, are you currently experiencing with the transition from the student role to the RN role?”; (b) “What could be done to help you feel more supported or integrated into the unit?”; (c) “What aspects of your work environment are most satisfying?”; and (d) “What aspects of your work environment are least satisfying?” The one open-ended question asks the respondent to “Please share any comments or concerns they have about their residency program.” Section V consists of 15 demographic questions.

Data Analysis

Data collected at the six-month period of graduate nurse responses were analyzed in this study. Using the split data file option in SPSS version 26 (SPSS Inc., Chicago, IL, USA), the total sample of 71,919 was split into three subsamples using random selection, yielding sample sizes of (a) EFA ($N= 23,489$), (b) CFA-1 ($N= 24,378$), and (c) CFA-2 ($N= 24,052$). Descriptive statistics were used to analyze the demographic information of

the sample. Prior to analysis, all items in Section II were labeled CF1 to CF24 and CF25a-f and all items in Section III were labeled CF26a-i. Negatively worded items were reverse coded prior to factor analysis. It was anticipated the Casey-Fink Survey would have subscales reflective of the latent variables or constructs measuring the phenomenon of role transition.

The researcher employed EFA as an exploratory or descriptive technique to determine the appropriate number of common factors and to uncover which measured variables were reasonable indicators of the various latent dimensions (e.g., by the size and differential magnitude of factor loadings; Brown, 2015). In EFA, all items freely loaded on factors and the solution was rotated to maximize the magnitude of primary loadings and minimize the magnitude of cross-loadings (Brown, 2015). Mathematically, EFA extracted maximum variance from the data set with each “factor” (Gray et al., 2017). The first factor was the linear combination of the survey items that maximized the variance of their factor scores. The second component was formed from residual variance. Subsequent factors were formed from the residual variances that had not yet been explained (Gray et al., 2017). There were two major types of rotation: orthogonal and oblique. In the orthogonal rotation, the factors were constrained to be uncorrelated (i.e., factors were oriented at 90-degree angles in multidimensional space). In the oblique rotation, the factors were allowed to intercorrelate (i.e., permit factor axis orientations of less than 90-degrees). The Varimax rotation is an orthogonal rotation method. Promax is an oblique rotation method.

Construct validity of the items in Sections II and III of the Casey-Fink Survey was tested through both PCA with Varimax rotation and principal axis factoring (PAF),

followed by Promax rotation with Kaiser normalization. Criteria for inclusion of an item on each factor included recommended minimum loadings of .40 and no cross-loadings greater than .15. The scree test involved examining the graph of the eigenvalues and looking for the natural bend or break point in the data where the curve flattened out (Costello & Osborne, 2005). The communality was a measure of the variable's shared variance (Polit, 2010).

Confirmatory factor analysis was conducted using AMOS statistical software (SPSS Inc., Chicago, IL, USA). In CFA, a model is specified, indicating which variables load on which factors and which factors are correlated. Confirmatory factor analysis is a form of structural equation modeling used to test hypothesized factor structures formulated from theory or suggested by prior empirical research (Brown, 2015). A measure of fit was indicated with this model. The model was analyzed and parameters estimated. The fit of the model was further evaluated. Model fit indices were examined to determine how well the model fit the data. The fit indices used to indicate a good fit included the comparative fit index (CFI) ≥ 0.95 , normed fit index (NFI) ≥ 0.95 , root mean square error of approximation (RMSEA) < 0.06 , and goodness of fit (GFI) ≥ 0.90 (Hu & Bentler, 1999; Tabachnick & Fidell, 2013). The chi-square test was reported but is not always a dependable fit index as a result of sample size issues (Polit & Yang, 2016). Because chi-square is sensitive to large sample sizes, its significance should not be ignored but it should be interpreted with caution (Kline, 2011).

Following EFA and CFA analyses, internal consistency reliability was computed for each sub-scale using SPSS and assessed using Cronbach's alpha. An important indicator of a scale's quality is its reliability coefficient—alpha. Nunnally (1978)

suggested a value of .70 as an acceptable lower bound for alpha. DeVellis (2017) suggested below .60 was unacceptable, between .60 and .65 was undesirable, between .65 and .70 was minimally acceptable, between .70 and .80 was respectable, between .80 and .90 was very good, and much above .90 one should consider shortening the scale.

The extent of missing data was evaluated on the EFA and both CFA samples. Missing values were identified during EFA analysis of items CF6 ($n = 404$, 1.8% of the EFA sample) and CF19 ($n = 423$; 1.8% of the EFA sample). Data values could be missing for any number of reasons including when study participants skipped over questions in a questionnaire, missed a data collection appointment, refused to continue in a study, or moved away from the study area (Polit, 2010). When values are missing, it means analyses are based on fewer study participants than were in the full study sample (Polit, 2010). The pattern of missing data needs to be addressed unless the amount missing is trivial, for example 1% missing (Polit, 2010). Missing data were not imputed as there was a small percentage (1.8%) of missing data and the conclusion was the missing data were completely at random.

Limitations of the Study

One limitation of this study was it was a self-report survey, which could lead to response bias. A potential threat to internal validity includes selection bias. All study participants were newly graduated nurses enrolled in the well-established, evidence-based Vizient/AACN (Vizient, 2018) 12-month Nurse Residency Program. Use of a convenience sample was a limitation of this study and social desirability was a potential response bias. Failure of participants to respond to all the questions on the survey instrument, especially with long questionnaires, could have also threatened the validity of

the instrument (Gray et al., 2017). These limitations could have affected the generalizability of the study findings to all populations of newly graduated nurses.

Summary

This chapter described the methodology used to answer the research question for this study. A discussion of secondary data analysis was included. The data collection section described the steps used to obtain the data for this study. The analysis of data section included a discussion of EFA and CFA statistical analyses conducted for this study, the importance for computing reliability coefficients on the data collected, and options for managing missing data. Additionally, this chapter included a description of the sample population, data management procedures, inclusion and exclusion criteria, ethical considerations, and study limitations.

CHAPTER IV

RESULTS

Chapter IV describes the results of this study. It begins with a description of the population sampled followed by a report of the results of the data analysis necessary to answer the research question that guided this study.

Description of the Sample

The analytic sample consisted of 71,919 survey responses collected from newly graduated nurses who participated in the Vizient/AACN (Vizient, 2018) 12-month Nurse Residency Program from 2008 to 2018. The data for this analysis were collected during a six-month time period. This final sample of 71,919 participants was randomly split into three samples called EFA ($N = 23,489$), CFA-1 ($N = 24,378$), and CFA-2 ($N = 24,052$). Descriptive statistics provided information related to the demographics of the sample. Participants in cohort years 2015 to 2018 comprised 71% ($N = 50,978$) of the total respondents. Table 2 provides a summary of participants for each cohort year.

The typical respondent was 26-years-old ($M = 26$ years) with a range of 45 years (20-65 years). Respondents were predominantly female (87.7%). The ethnicity of the sample included African American (8.1%), American Indian (0.2%), Asian (7.3%), Caucasian (74%), Hispanic (5.6%), Pacific Islander (0.2%), two or more races (1.5%), and Other (2.6%). Most of the respondents reported receiving a Bachelor of Science in Nursing degree (61.6%) and a majority did not have a previous non-nursing degree (63.4%). Demographic characteristics across all three randomly selected samples used

for the factor analysis were similar. Demographics for the total sample, EFA sample, CFA-1 sample, and CFA-2 sample can be found in Table 3.

Table 2

Participants by Cohort Year

Cohort Year	<i>N</i>	%	Valid %	Cumulative %
2008	1,842	2.6	2.6	2.6
2009	2,006	2.8	2.8	5.4
2010	2,422	3.4	3.4	8.7
2011	2,322	3.2	3.2	11.9
2012	2,905	4.0	4.0	16.0
2013	3,377	4.7	4.7	20.7
2014	6,067	8.4	8.4	29.1
2015	8,551	11.9	11.9	41.0
2016	11,354	15.8	15.8	56.8
2017	14,992	20.8	20.8	77.6
2018	16,081	22.4	22.4	100.0
Total	71,919	100.0	100.0	

Table 3

Demographic Characteristics of the Six-Month Samples

Characteristic	Total 6-month Sample (N= 71,919) %	EFA Sample (n = 23,489) %	CFA-1 Sample (n = 24,378) %	CFA-2 Sample (n= 24,052) %
Age (years)	<i>M</i> = 26.4 <i>SD</i> ± 6.1	<i>M</i> = 26.4 <i>SD</i> ±6.2	<i>M</i> = 26.4 <i>SD</i> ±6.2	<i>M</i> = 26.4 <i>SD</i> ±6.0
Gender				
Female	87.7	87.8	87.6	87.8
Male	12.1	12.1	12.2	12.1
No response	0.2	0.1	0.2	0.2
Ethnicity				
African American	8.7	8.7	8.7	8.7
American Indian	0.2	0.2	0.2	0.2
Asian	7.3	7.2	7.3	7.3
Caucasian	74.0	73.7	74.2	74.0
Hispanic or Latino	5.6	5.7	5.4	5.6
Pacific Islander	0.2	0.2	0.2	0.2
Unknown	2.6	2.7	2.5	2.5
No response	1.5	1.5	1.5	1.5
Nursing Degree Received				
Diploma	1.1	1.1	1.1	1.0
Associate Degree	20.6	20.6	20.5	20.6
BSN	61.6	61.6	61.7	61.5
Accelerated BSN	14.3	14.2	14.2	14.5
Master's	2.0	2.0	2.0	2.1
No response	0.4	0.4	0.4	0.3
Previous Non-Nursing Degree				
Associate Degree	7.1	7.1	7.1	7.1
Baccalaureate	27.3	27.1	27.0	27.7
Master's	1.9	1.9	2.0	1.9
Doctorate	0.3	0.3	0.3	0.3
No previous degree	63.4	63.6	63.5	62.9

Note. Percentages may not add up to 100% due to rounding error.

The largest percentage of the respondents worked on Adult Medical/Surgical Units (26%) followed by Adult Medical/Surgical Intensive Care Units (12.3%). The Emergency Department employed an average of 6.5% of respondents, Telemetry Units employed 7.5%, and 10.6% entered no response or selected “other” as their unit of

employment. Responses to clinical unit of employment across all three randomly selected samples were similar. The clinical units of employment for the total six-month sample, the EFA sample, the CFA-1 sample, and the CFA-2 sample can be found in Table 4.

Table 4

Clinical Unit of Employment for the Six-Month Samples

Clinical Unit of Employment	Total 6-month Sample (N= 71,919) %	EFA Sample (n = 23,489) %	CFA-1 Sample (n = 24,378) %	CFA-2 Sample (n= 24,052) %
Ambulatory	0.3	0.3	0.3	0.2
Burn care	0.3	0.3	0.2	0.3
Cardiac intensive care unit	2.5	2.5	2.5	2.6
Emergency department	6.6	6.5	6.7	6.6
Float pool	1.4	1.5	1.4	1.5
Home care	0.0	0.0	0.0	0.0
Hospice/palliative care	0.0	0.0	0.0	0.0
Labor & delivery	3.1	3.3	2.9	3.1
Medical/Surgical	26.3	26.6	26.3	25.9
Medical/Surgical intensive care unit	12.3	12.1	12.3	12.6
Medical/Surgical intermediate	2.2	2.1	2.2	2.1
Mother/Baby	1.4	1.4	1.3	1.4
Neonatal intensive care unit	3.3	3.4	3.2	3.3
Neurosurgical intensive care unit	1.7	1.6	1.7	1.7
Newborn nursery	0.0	0.0	0.0	0.0
No response	2.3	2.4	2.3	2.3
Oncology	5.4	5.3	5.4	5.5
Orthopedic	1.5	1.5	1.6	1.4
Other	8.3	8.1	8.4	8.5
Pediatric emergency department	0.2	0.2	0.2	0.2
Pediatric intensive care unit	2.3	2.4	2.2	2.2
Pediatric	3.5	3.4	3.6	3.5
Pediatric intermediate	0.2	0.2	0.3	0.2
Pediatric oncology	0.4	0.4	0.3	0.4
Pediatric/Adolescent	0.1	0.1	0.1	0.2
Psychiatric				
Perioperative (operating room)	3.2	3.3	3.2	3.2
Perioperative (post-anesthesia care unit)	0.5	0.5	0.5	0.5
Psychiatric	1.0	1.0	1.0	1.0
Rehabilitation	0.5	0.4	0.6	0.5
Skilled nursing facility	0.0	0.0	0.0	0.0
Telemetry	7.5	7.5	7.3	7.7
Transplant	1.5	1.4	1.5	1.6

Note. Percentages may not add up to 100% due to rounding error.

Analysis of Variables

Each of the 24 items in Section II of the Casey-Fink Survey was measured using a 4-point Likert response scale. Means for the 24 items for each sample in the six-month time period are presented in Table 5. The item means across all three randomly selected samples were similar. Responses to the six questions asking for causes of stress are listed in Table 6. These items were measured using a yes/no (binary) format.

Table 5

Item Means for Section II for Each Sample in the Study

Item	EFA Sample <i>n</i> = 23,489 <i>M</i> (<i>SD</i>)	CFA-1 Sample <i>n</i> = 24,378 <i>M</i> (<i>SD</i>)	CFA-2 sample <i>n</i> = 24,052 <i>M</i> (<i>SD</i>)	Total Sample <i>N</i> = 71,919 <i>M</i> (<i>SD</i>)
CF1 I feel confident communicating with Physicians.	3.21 (.527)	3.21 (.537)	3.21 (.531)	3.21 (.532)
CF2 I am comfortable knowing what to do for a dying patient.	2.65 (.717)	2.65 (.721)	2.64 (.725)	2.65 (.721)
CF3 I feel comfortable delegating tasks to the Nursing Assistant.	3.19 (.584)	3.19 (.587)	3.19 (.589)	3.19 (.587)
CF4 I feel at ease asking for help from other RNs on the unit	3.52 (.569)	3.52 (.567)	3.53 (.568)	3.52 (.568)
CF5 I am having difficulty prioritizing patient care needs	3.05 (.634)	3.04 (.642)	3.05 (.628)	3.04 (.635)
CF6 I feel my preceptor provides encouragement and feedback about my work.	3.43 (.639)	3.43 (.643)	3.42 (.639)	3.43 (.640)
CF7 I feel staff is available to me during new situations and procedures.	3.34 (.605)	3.33 (.616)	3.34 (.609)	3.34 (.610)
CF8 I feel overwhelmed by my patient care responsibilities and workload	2.71 (.718)	2.72 (.719)	2.72 (.714 0	2.72 (.717)
CF9 I feel supported by the nurses on my unit.	3.42 (.577)	3.42 (.582)	3.43 (.584)	3.42 (.581)
CF10 I have opportunities to practice skills and procedures more than once.	3.18 (.577)	3.19 (.584)	3.18 (.575)	3.18 (.579)
CF11 I feel comfortable communicating with patients and their families.	3.34 (.524)	3.34 (.525)	3.34 (.520)	3.34 (.523)
CF12 I am able to complete my patient care assignment on time.	3.13 (.587)	3.13 (.590)	3.12 (.582)	3.13 (.586)
CF13 I feel the expectations of me in this job are realistic.	2.99 (.639)	3.00 (.635)	2.99 (.627)	2.99 (.634)
CF14 I feel prepared to complete my job responsibilities.	3.14 (.504)	3.15 (.503)	3.14 (.496)	3.15 (.501)
CF15 I feel comfortable making suggestions for changes to the nursing plan of care.	2.98 (.594)	2.98 (.596)	2.97 (.594)	2.98 (.595)
CF16 I am having difficulty organizing patient care needs	3.04 (.617)	3.05 (.620)	3.05 (.607)	3.05 (.615)
CF17 I feel I may harm a patient due to my lack of knowledge and experience	3.09 (.700)	3.09 (.702)	3.09 (.690)	3.09 (.697)
CF18 There are positive role models for me to observe on my unit.	3.48 (.552)	3.48 (.548)	3.48 (.551)	3.48 (.550)
CF19 My preceptor is helping me to develop confidence in my practice	3.41 (.647)	3.40 (.653)	3.41 (.645)	3.41 (.648)
CF20 I am supported by my family/friends.	3.57 (.531)	3.57 (.528)	3.58 (.529)	3.57 (.529)
CF21 I am satisfied with my chosen nursing specialty.	3.22 (.712)	3.22 (.718)	3.22 (.716)	3.22 (.716)
CF22 I feel my work is exciting and challenging.	3.28 (.609)	3.28 (.620)	3.29 (.614)	3.28 (.614)
CF23 I feel my manager provides encouragement and feedback about my work.	2.98 (.798)	2.98 (.797)	2.98 (.795)	2.98 (.797)
CF24 I am experiencing stress in my personal life.	2.45 (.812)	2.44 (.816)	2.46 (.811)	2.45 (.813)

Table 6

Causes of Stress

Stressors	EFA		CFA-1		CFA-2		Total	
	Yes %	No %	Yes %	No %	Yes %	No %	Yes %	No %
CF25a Financial Stress	23.3	76.7	23.2	76.8	24.0	76.0	23.5	76.5
CF25b Child Care	5.3	94.7	5.2	94.8	4.9	95.1	5.1	94.9
CF25c Student Loans	21.9	78.1	21.9	78.1	22.7	77.3	22.2	77.8
CF25d Living Situation	11.5	88.5	11.0	89.0	11.4	88.6	11.3	88.7
CF25e Personal Relationship	21.6	78.4	21.0	79.0	21.6	78.4	21.4	78.6
CF25f Job Performance	12.3	87.7	12.1	87.9	12.6	87.4	12.3	87.7

Each of the nine items in Section III of the survey was measured using a 5-point Likert response scale. The means for items in the EFA sample, CFA-1 sample, and CFA-2 sample in Section III of the survey can be found in Table 7. The item means across all three randomly selected samples were similar.

Table 7

Item Means for Section III for Each Sample in the Study

Item	EFA Sample <i>n</i> = 23,489 <i>M</i> (<i>SD</i>)	CFA-1 Sample <i>n</i> = 24,378 <i>M</i> (<i>SD</i>)	CFA-2 Sample <i>n</i> = 24,052 <i>M</i> (<i>SD</i>)	Total Sample <i>N</i> = 71,919 <i>M</i> (<i>SD</i>)
CF26a Salary	3.55 (1.03)	3.54 (1.03)	3.54 (1.03)	3.54 (1.03)
CF26b Vacation	3.56 (1.00)	3.57 (1.00)	3.57 (.999)	3.57 (1.03)
CF26c Benefits Package	3.83 (.89)	3.83 (.89)	3.82 (.895)	3.83 (.894)
CF26d Hours Worked	3.92 (.882)	3.92 (.877)	3.91 (.881)	3.92 (.880)
CF26e Weekends off/month	3.42 (1.19)	3.43 (1.18)	3.40 (1.18)	3.41 (1.19)
CF26f Work Responsibilities	3.77 (.852)	3.78 (.855)	3.77 (.842)	3.77 (.850)
CF26g Opportunities for Career Advancement	3.84 (.851)	3.84 (.851)	3.83 (.849)	3.84 (.850)
CF26h Encouragement and Feedback	3.80 (.945)	3.79 (.95)	3.79 (.951)	3.79 (.949)
CF26i Opportunities for Choosing shifts Worked	3.62 (1.10)	3.63 (1.10)	3.62 (1.10)	3.62 (1.10)

Exploratory Factor Analysis

To answer the research question, “What are the underlying constructs measured by the items in Sections II and III on the Casey-Fink Survey in newly graduated nurses who participated in the one-year Vizient/AACN TM Nurse Residency Program,” an exploratory factor analysis was conducted on the EFA sample of 23,489 graduate nurse responses at the six-month time period.

A PCA approach was selected as the initial method of factor extraction. Principal components analysis is used when the researcher has no pre-existing knowledge about the factors that might explain the inter-relationships between a set of variables (Pett et al., 2003). A Varimax rotation (orthogonal rotation) was chosen to maximize the variance of

the loadings within factors and across variables (Polit, 2010). The Kaiser-Meyer-Olkin measure and the Bartlett test of sphericity were used to evaluate appropriateness of factor analysis. The Kaiser-Meyer-Olkin measure of sampling adequacy could range from 0 to 1 (Polit, 2010). The closer the value was to 1, the better the prospects for factor analysis (Polit, 2010).

The 24 items that comprised Section II of the Casey-Fink Survey were analyzed in the initial analysis using PCA and Varimax (orthogonal) rotation to identify the number of factors to extract. All item responses were reported using a 4-point Likert scale format. A six-factor solution was found with eigenvalues exceeding 1, accounting for 60% of the variance in total scores. The Kaiser-Meyer-Olkin index of sampling adequacy was .893, which is considered “meritorious” by Kaiser (1974), and the Bartlett’s test of sphericity was statistically significant ($p < .001$). Significance on the Bartlett’s test of sphericity supported further evaluation of the factorability of the data (Polit, 2010). The communality scores ranged from .323 to .926. Retention of items on factors was based on the recommended minimum loading of .40 and cross-loadings of greater than .15. In this initial analysis, all items loaded on two or more of the six factors; three items (CF8, CF12, and CF14) had cross-loadings greater than .15 and were, therefore, eliminated. Item CF10 (“I have opportunities to practice skills and procedures more than once”) and CF24 (“I am experiencing stress in my personal life”) failed to load on any factor and were also eliminated. Factor 1 was comprised of five items with factor loadings that ranged from .59 to .76. Factor 2 was comprised of five items with factor loadings that ranged from .50 to .79. Factor 3 was comprised of three items with factor loadings that ranged from .63 to .85. Factor 4 was comprised of three items with factor

loadings that ranged from .45 to .77. Factor 5 was comprised of three items with factor loadings that ranged from .44 to .78. Factor 6 was comprised of two items with factor loadings of .95 to .96. The two items in Factor 6 described the preceptor role. Inspection of the scree plot revealed a break in the slope at Factors 5 and 6, which directed subsequent analysis.

A second factor analysis using a principal axis factoring (PAF) approach was conducted on the 24 items of Section II of the Casey-Fink Survey. A principal axis factoring approach was selected to identify the latent constructs underlying the scale. Promax (oblique) rotation was selected. An oblique rotation (Promax) was chosen because it was hypothesized the underlying dimensions of the constructs in Section II of the Casey-Fink Survey would be correlated (Polit, 2010). Promax rotation allowed for correlations among the factors and was extremely useful for larger data sets (Pett et al., 2003). The Kaiser-Meyer-Olkin index of sampling adequacy was .893 and the Bartlett's test of sphericity was statistically significant ($p < .001$). A six-factor solution was found with eigenvalues exceeding 1, accounting for 54% of the variation in total scores. The factor pattern matrix was used to determine the factor scores. Criteria for inclusion of an item on a factor were a minimum of .40 and no cross-loading greater than .15. Factor 1 was comprised of five items with factor loadings of .46 to .89. Factor 2 was comprised of five items with factor loadings of .49 to .67. Factor 3 was comprised of three items with factor loadings of .54 to .77. Factor 4 was comprised of two items with factor loadings of .93 to .94. Factor 5 was comprised of two items with factor loadings of .81 to .84. Factor 6 was comprised of three items with factor loadings of .52 to .90. Both Factors 4 and 5 included only two items, below the ideal number of items per factor (Polit & Beck,

2012). Factor 3 had three items that loaded together although they were the negatively worded items that had been recoded. This finding suggested similar items might hold together due to the negative wording or it might be that those items were measuring the same construct. In this six-factor solution, more items cross-loaded onto multiple factors and did not present “clean loadings.” Ideally, the researcher would like an item to load significantly on a single factor (Pett et al., 2003). Item CF8 was eliminated due to cross-loadings of $>.15$. Items CF10 (“I have opportunities to practice skills and procedures more than once”), CF23 (“I feel my manager provides encouragement and feedback about my work”), and CF24 (“I am experiencing stress in my personal life”) did not load on any of the factors due to their low factor loadings. Thus, these three items were eliminated.

Inspection of the scree plot revealed a break in the slope at Factors 4 and 6. Since the goal was to obtain the most parsimonious solution, it was decided to conduct a parallel analysis to decide how many factors to extract and retain. Parallel analysis is a statistical method used to compare eigenvalues generated from the data mix to the eigenvalues generated from the Monte-Carlo simulated matrix created from random data of the same size (Tran & Formann, 2009). Factors are retained when actual eigenvalues surpass random ordered eigenvalues (Williams, Brown, & Onsman, 2010). The random ordered eigenvalue variance for a five-factor solution using parallel analysis was 1.140 and the actual PCA eigenvalue variance for a five-factor solution was 1.354, which surpassed the random ordered eigenvalues. It was decided to conduct further exploratory factor analysis specifying five factors to be extracted.

The stress item (CF24) had consistently low loadings ($>.40$) on all factors and was therefore a good candidate for deletion. Conceptually, graduate nurses experience stress when transitioning from the student to professional nurse role (Casey et al., 2004), so it was decided to add the six responses to specific items participants identified as “causing them stress” into the exploratory factor analysis. The six items added were financial stress (labeled CF25a), child care stress (CF25b), student loan stress (labeled 25c), living situation stress (labeled CF25d), personal relationship stress (labeled CF25e), and job performance stress (labeled CF25f). These six items used a binary response format and were recorded as Yes and No responses in the data set. These items were re-coded as 1= Yes and 2= No.

The third EFA used principal axis factoring (PAF) extraction with Varimax (orthogonal) rotation, specifying five factors to be extracted, was conducted on the 24 items in Section II of the survey including the additional CF25a-f stress items. Principal axis factoring was selected for all subsequent EFA extraction methods as this approach looked at common variance between the items. The Kaiser-Meyer-Olkin index of sampling adequacy was .893, and Bartlett’s test of sphericity was statistically significant ($p < .000$). A five-factor solution was specified and the cumulative variance was 49.7% of total scores. The rotated factor matrix was used to determine the factor scores. Criteria for inclusion of an item on a factor were a minimum of .40 and no cross-loading greater than .15. Factor 1 was comprised of eight items with factor loadings of .42 to .74. This factor was labeled Social Support as it reflected perceptions of support received from coworkers on the unit of employment. Factor 2 was comprised of seven items with factor loadings of .42 to .66. This factor was labeled Role Confidence as these items

related to the professional role responsibilities a new nurse is learning. Factor 3 was comprised of five items with factor loadings of .47 to .76. This factor was labeled Stress as the items reflected personal stressors new nurses encountered in the first year of practice. All items except CF25b (child care stress) and CF25f (job performance stress), had moderate loadings on the stress factor. Factor 4 was comprised of four items with factor loadings of .46 to .72. These items included the four negatively worded items that related to organization/prioritization of patient care needs. This factor was labeled ability to Organize/Prioritize care. Factor 5 was comprised of two items with factor loadings of .87 to .88. This factor was labeled Professional Socialization as both items suggested the importance of the preceptor in guiding and socializing the new nurse into their role on the unit of employment during role transition. It was noticed the items that loaded onto all five factors did not have as many cross-loadings as in previous extractions. Items CF2 (“I am comfortable knowing what to do for a dying patient”) and CF10 (“I have opportunities to practice skills and procedures more than once”) failed to load on any factor and were therefore eliminated.

The 30-item scale was then tested with the EFA sample ($n = 23,489$) using PAF extraction and Promax (oblique) rotation and specified five factors to extract. This analysis yielded five conceptually clear factors and resulted in the elimination of four of the total items. These five factors explained 49.7% of the extracted common variance. Using the pattern matrix, a minimum factor loading of .40 was used as a criterion for each retained item and at least a difference of .15 between the primary loading and any secondary loading for an item. Factor 1 (support) was comprised of seven items with factor loadings of .53 to .85. Factor 2 (role confidence) was comprised eight items with

factor loadings of .43 to .72. Factor 3 (stress) was comprised of five items with factor loadings of .48 to .76. Item CF25b (child care stress) and CF25f (job performance stress) were eliminated due to low loadings of .233 and .355. Factor 4 (organize/prioritize care) was comprised of four items and had factor loadings of .44 to .78. Factor 5 (professional socialization) was comprised of two items with factor loadings of .92 to .93. It was decided to keep this factor with only two items as the preceptor role is vital to a new nurse's role transition (Casey et al., 2004). Item CF10 ("I have opportunities to practice skills and procedures more than once") and CF23 ("I feel my manager provides encouragement and feedback about my work") were eliminated as they both loaded low on two factors. This five-factor EFA solution of 26 items provided the basis for a hypothesized model that was used for the CFA-1 testing. Table 8 presents the final five-factor item loadings used in the CFA testing. This final solution resulted in 26 items for the confirmatory factor analysis.

A summary of the EFA models, methods for testing the items in Section II of the survey, and results obtained are summarized in Table 9.

Table 8

Exploratory Factor Analysis Five-Factor Loadings for Items in Section II

Pattern Matrix	Factor				
	1	2	3	4	5
CF9 I feel supported by the nurses on my unit.	.848	-.140			
CF18 There are positive role models for me to observe on my unit.	.841	-.182			
CF7 I feel staff is available to me during new situations and procedures.	.736				
CF20 I am supported by my family/friends.	.598				
CF4 I feel at ease asking for help from other RNs on the unit	.584				
CF22 I feel my work is exciting and challenging.	.557	.167		-.106	
CF21 I am satisfied with my chosen nursing specialty.	.530	.169			
CF23 I feel my manager provides encouragement and feedback about my work.	.381	.169		-.144	
CF14 I feel prepared to complete my job responsibilities.		.716			
CF15 I feel comfortable making suggestions for changes to the nursing plan of care.		.690			
CF12 I am able to complete my patient care assignment on time.		.579		.153	
CF1 I feel confident communicating with Physicians.		.565			
CF13 I feel the expectations of me in this job are realistic.	.173	.523			
CF2 I am comfortable knowing what to do for a dying patient.	-.157	.499		-.101	
CF11 I feel comfortable communicating with patients and their families.	.171	.494			
CF3 I feel comfortable delegating tasks to the Nursing Assistant.		.430			
CF10 I have opportunities to practice skills and procedures more than once.	.334	.358			
CF24 I am experiencing stress in my personal life.			.755		
CF25a Financial Stress			.691		
CF25c Student Loan Stress			.594		
CF25e Personal relationship Stress			.546		
CF25d Living Situation Stress			.479		
CF25f Job Performance Stress			.355	-.112	
CF25b Child Care Stress			.233		
CF16 I am having difficulty organizing patient care needs				.784	
CF5 I am having difficulty prioritizing patient care needs				.753	
CF17 I feel I may harm a patient due to my lack of knowledge and experience		.104		.527	
CF8 I feel overwhelmed by my patient care responsibilities and workload		.149		.435	
CF6 I feel my preceptor provides encouragement and feedback about my work.					.927
CF19 My preceptor is helping me to develop confidence in my practice					.917

Extraction Method: Principal Axis Factoring.

Rotation Method: Promax with Kaiser Normalization.

Bolded entries represent items retained on each factor.

Table 9

Summary of Factor Analysis Models for Section II

Model	Methods	Results
Model I	Principal component analysis, Varimax rotation, on 24 items	6-factor solution explained 60% of total variance with item loadings ranging from .32 -.93
Model II	Principal axis factoring, Promax rotation, on 24 items	6-factor solution explained 54% of total variance with item loadings ranging from .46 -.94
Model III	Principal axis factoring, Varimax rotation, on 30 items	5-factor solution explained 49.7% of total variance with item loadings ranging from .40 -.88
Model IV	Principal axis factoring, Promax rotation, on 30 items	5-factor solution explained 49.7% of total variance with item loadings ranging from .43 -.93

A separate exploratory factor analysis using the nine items in section III of the Casey-Fink Survey, was conducted with the EFA sample ($n = 23,489$). Principal component analysis extraction and Varimax (orthogonal) rotation was the approach for this analysis. Principal component analysis was used as this was exploratory in nature and no previous factor analysis had been reported on this scale. All item responses were reported using a 5-point Likert scale format. A two-factor solution was found with eigenvalues exceeding 1, accounting for 53.14% of the variation in total scores. The Kaiser-Meyer-Olkin measure of sampling adequacy was .87 and the Bartlett's test of sphericity was statistically significant ($p < .000$), which confirmed the appropriateness of the factor analysis. Criteria for inclusion of an item on a factor were a minimum loading of .40 and no cross-loading greater than .15 (Kline, 2011). The scree plot showed a clear "elbow" bend at two factors, which further supported keeping these two factors. Factor 1 was comprised of six items (CF26d-i) with factor loadings of .57 to .73 and was labeled Internal Benefits as these items reflected personal aspects of job satisfaction including

benefits offered in the workplace or unit of employment. Factor 2 was comprised of three items (CF26 a-c) with factor loadings of .69 to .77 and was labeled External Benefits. These three items reflected perceptions of external benefits = the hiring organization provided to graduate nurses. Factor loadings for the nine items in Section III of the survey are presented in Table 10. These nine items were used for the confirmatory factor analysis testing.

Table 10

Exploratory Factor Analysis Loadings for Items in Section III

Rotated Component Matrix	Component	
	1	2
CF26h Encouragement and feedback	.735	.135
CF26i Opportunity for choosing shifts worked	.705	
CF26f Work responsibilities	.663	.260
CF26g Opportunities for career advancement	.660	.299
CF26e Weekends off per month	.593	.170
CF26d Hours worked	.570	.388
CF26c Benefits package	.182	.775
CF26a Salary	.139	.768
CF26b Vacation	.314	.698

Extraction Method: Principal Component Analysis.

Rotation Method: Varimax with Kaiser Normalization.

Bolded entries represent items that were retained on each factor.

Confirmatory Factor Analysis

A confirmatory factor analysis (CFA) approach was used to evaluate the extent to which the 26 items fit the specified five-factor structure model that resulted from the EFA conducted on Section II of the survey. It was hypothesized these 26 items belonged to the five-factor model and CFA helped determine if there was a good-fit with the specified factors (see Figure 1). Fit refers to how close the observed data match the relationships specified in the hypothesized model (Brown, 2015). Data used for the CFA-1 testing were a separate, random sample of graduate nurse responses at the six-month time period ($n= 24,378$). The measurement model was created in AMOS to reflect the relationships between the variables and the factors. The two directional arrows shown in this model indicated there might be a correlation between the five factors (see Figure 2).

Assessment of model fit was evaluated using multiple fit indices since each index provided information on different aspects of model fit (Polit & Yang, 2016). The fit statistics used as a threshold for acceptance were GFI, NFI, CFI, and RMSEA.

Acceptable model fit was defined by the following criteria: GFI (≥ 0.90), NFI (≥ 0.95), RMSEA (≤ 0.06), and CFI (≥ 0.95 ; Hu & Bentler, 1999; Tabachnick & Fidell, 2013).

Figure 1. Confirmatory factor analysis-1 five-factor model for 26 items in section II.

Figure 2. Confirmatory factor analysis-1 model for items in section III.

The model was drawn in AMOS and the parameters and analysis properties were set. Maximum likelihood estimation was not possible due to the number of missing responses on the professional socialization items. The estimate means and intercepts were selected in the analysis properties to account for the missing data. The model fit output for the baseline CFA-1 (Model A) indicated a poor fit to the data (see Table 11). The regression weights were reviewed for any problematic items. Items with the weakest coefficients were removed to improve model fit. Weak standardized factor loading coefficients were identified on CF2 (.762), CF26d (.620), and CF20 (.710). Based on these values, these three items were removed. The CF24 (stress question) showed a negative loading of -2.773 and was kept in the model as it was hypothesized stress was an

important contributor to the content of the scale. The revised model was re-run as Model B. The model fit output for Model B is shown in Table 11. Based on these results and the persistent low and negative loadings of the items on the stress factor, CF24 and CF25a-f were removed to improve model fit. This four-factor model was then labeled Model C. The model fit output for Model C is shown in Table 11. The model fit did not improve. The regression weights were reviewed and it was decided to remove the socialization factor (CF 6 and CF19) as the missing values impeded the ability of AMOS to run modification indices. This three-factor model was labeled Model D. The model fit output for Model D is shown in Table 11. The modification indices could then be reviewed to identify pairs of items that had high covariance. Items e6 to e1 and items e7 to e6 demonstrated large modification indices and were co-varied to improve model fit. Normally, one would not want error terms to have high covariance. The placement of two items (CF21 and CF22) related to professional satisfaction and might not match the wording of item CF9, which related to support provided by nurses. This model was labeled Model E. The model fit output for Model E is shown in Table 11. Table 11 provides a summary of the model fit output and action taken in CFA-1.

Table 11

Confirmatory Factor Analysis-1 Model Output and Actions Taken

Model	RMSEA	CFI	NFI	Action Taken
A	.068	.864	.863	Baseline
B	.073	.869	.868	CF2, 25d, and CF20 removed
C	.085	.870	.869	Stress factor (CF24, 25a-e) removed
D	.095	.837	.836	Socialization factor (CF9, 16) removed
E	.072	.908	.970	Co-vary e6 to e1 and e6 to e7

Note. RMSEA = Root mean square error of approximation, CFI = Comparative fit index, NFI = Normed fit index.

These CFA-1 results suggested a poor fit of the data despite the removal of weak items and the addition of co-variances between error terms on the same factor where large modification indices was observed. Removal of the stress and socialization factors due to negative correlations and modification estimation could not be completed due to missing data. It was determined there were too many problems with the initial EFA five-solution model structure, which resulted in too many modifications to achieve good model fit.

A CFA-1 was also conducted on the nine items in section III (Job Satisfaction) of the Casey-Fink Survey to test the extent to which the data fit the specified two-factor structure model that resulted from the EFA conducted on Section III of the survey. The model was drawn in AMOS (see aforementioned Figure 2) and the run parameters and analysis properties were set. Maximum likelihood estimation was possibly due to no

missing responses in the data. Model fit output using the CFA-1 sample ($n = 24,378$) was CFI = .925, NFI = .925, and RMSEA = .084, which suggested an adequate model fit. It was concluded the two factors (External Benefits and Internal Benefits) of nine items had desirable CFA results, suggesting good fit with the data. Figure 2 presented the CFA-1 model for items in Section III.

After reviewing role transition theory and results of the CFA-1 analysis for Sections II and III of the Casey-Fink Survey, this researcher decided to combine the 24 items in Section II that related to transition into the professional nurse role and the nine items in Section III, which related to job satisfaction, and conducted another exploratory factor analysis on the combined 33 items of the survey. It was hypothesized role transition constructs derived from theory might have been included in both Sections II and III in the survey. The six questions (CF25a-f) that asked about the causes of stress graduate nurses reported were not included in this final EFA analysis. The binary responses to these six questions were problematic in testing the CFA-1 model and are discussed as a measurement issue for this survey. A parallel analysis was re-run as the number of items included in the analysis was changed.

The final EFA—using PAF extraction, Promax rotation, and specifying five factors to be extracted—was conducted on the combined 33 items in both Sections II and III of the survey. The analysis used the EFA sample ($n = 23,489$) graduate nurse responses at the six-month time period. Stress items 25a-f were not included as the binary response format was found to be problematic. This final solution resulted in the Kaiser-Meyer-Olkin index of sampling adequacy of .915, which was considered “marvelous” by Kaiser (1974), and the Bartlett’s test of sphericity was statistically

significant ($p < .000$; see Table 12). The communality scores ranged from .140 to .783 (see Table 13).

Table 12

Results of Kaiser-Meyer-Olkin Index of Sampling Adequacy and Bartlett's Test of Sphericity

Final Solution		
Kaiser-Meyer-Olkin Measure of Sampling Adequacy.		.915
Bartlett's Test of Sphericity	Approx. Chi-Square	294987.631
	<i>df</i>	528
	Sig.	.000

Table 13

Communalities Scores

Communalities	Initial score
CF1 I feel confident communicating with Physicians.	.307
CF2 I am comfortable knowing what to do for a dying patient.	.140
CF3 I feel comfortable delegating tasks to the Nursing Assistant.	.276
CF4 I feel at ease asking for help from other RNs on the unit	.417
CF6 I feel my preceptor provides encouragement and feedback about my work.	.782
CF7 I feel staff is available to me during new situations and procedures.	.465
CF9 I feel supported by the nurses on my unit.	.553
CF10 I have opportunities to practice skills and procedures more than once.	.369
CF11 I feel comfortable communicating with patients and their families.	.394
CF12 I am able to complete my patient care assignment on time.	.452
CF13 I feel the expectations of me in this job are realistic.	.549
CF14 I feel prepared to complete my job responsibilities.	.550
CF15 I feel comfortable making suggestions for changes to the nursing plan of care.	.333
CF18 There are positive role models for me to observe on my unit.	.487
CF19 My preceptor is helping me to develop confidence in my practice	.783
CF20 I am supported by my family/friends.	.351
CF21 I am satisfied with my chosen nursing specialty.	.569
CF22 I feel my work is exciting and challenging.	.559
CF23 I feel my manager provides encouragement and feedback about my work.	.421
CF24 I am experiencing stress in my personal life.	.086
CF5 I am having difficulty prioritizing patient care needs.	.339
CF8 I feel overwhelmed by my patient care responsibilities and workload.	.405
CF16 I am having difficulty organizing patient care needs.	.388
CF17 I feel I may harm a patient due to lack of knowledge and experience.	.300
CF26a Salary	.270
CF26b Vacation	.363
CF26c Benefits package	.311
CF26d Hours worked	.349
CF26e Weekends off per month	.254
CF26f Work responsibilities	.515
CF26g Opportunities for career advancement	.419
CF26h Encouragement and feedback	.549
CF26i Opportunity for choosing shifts worked	.308

Extraction Method: Principal Axis Factoring.

The final EFA analysis resulted in five factors that explained 49.49% of the extracted common variance. Figure 3 shows the scree plot of eigenvalues used to determine the number of significant factors.

Figure 3. Scree plot for final exploratory factor analysis.

The pattern matrix was examined and a minimum factor loading of .40 was used as a criterion for each retained item and at least a difference of .15 between the primary loading and any secondary loading for an item. Factor 1 was comprised of nine items (CF26a-i) with factor loadings of .52 to .68. Factor 2 was comprised of five items (CF4, CF7, CF9, CF18, and CF20) with factor loadings of .60 to .84. Factor 3 was comprised of eight items (CF1, CF2, CF3, CF11, CF12, CF13, CF14, and CF15) with factor loadings of .60 to .84. Factor 4 was comprised of four items (CF5, CF8, CF16, and CF17) with factor loadings of .46 to .75. Factor 5 was comprised of two items (CF6 and CF19) with factor loadings of .94 to .95. This factor with only two items was retained in the model as the preceptor role is vital to a new nurse's role transition (Casey et al., 2004). Items CF10 ("I have opportunities to practice skills and procedures more than once"), CF21 ("I am satisfied with my chosen nursing specialty"), CF22 ("I feel my work is exciting and challenging"), CF23 ("I feel my manager provides encouragement and feedback about my work"), and CF24 ("I am experiencing stress in my personal life")

were eliminated as they all loaded less than .40 and had loadings on multiple factors.

This five-factor EFA solution with 28 items provided the basis for a second CFA model testing (see Table 14 for the final five-factor solution).

A second confirmatory factor analysis was conducted on the 28 items from Sections II and III of the survey to test the fit to the data (see Figure 4 for the model used for CFA-1 analysis).

Table 14

Pattern Matrix for the Final Five-Factor Solution

Items	Factor				
	1	2	3	4	5
CF26b Vacation	.684				
CF26d Hours worked	.641				
CF26g Opportunities for career advancement	.639				
CF26c Benefits package	.604				
CF26f Work responsibilities	.602		.215		
CF26a Salary	.595				
CF26e Weekends off per month	.559				
CF26i Opportunity for choosing shifts worked	.541				
CF26h Encouragement and feedback	.518				
CF23 I feel my manager provides encouragement and feedback about my work.	.311	.207			
CF9 I feel supported by the nurses on my unit.	.845				
CF18 There are positive role models for me to observe on my unit.	.814				
CF7 I feel staff is available to me during new situations and procedures.	.699				
CF4 I feel at ease asking for help from other RNs on the unit	.668				
CF20 I am supported by my family/friends.	.599				
CF22 I feel my work is exciting and challenging.	.215	.356	.209		
CF21 I am satisfied with my chosen nursing specialty.	.257	.305	.211		
CF14 I feel prepared to complete my job responsibilities.			.772		
CF15 I feel comfortable making suggestions for changes to the nursing plan of care.			.706		
CF12 I am able to complete my patient care assignment on time.			.643		
CF13 I feel the expectations of me in this job are realistic.	.238		.591		
CF1 I feel confident communicating with Physicians.			.587		
CF11 I feel comfortable communicating with patients and their families.	.224		.535		
CF2 I am comfortable knowing what to do for a dying patient.			.493		
CF3 I feel comfortable delegating tasks to the Nursing Assistant.			.452		
CF10 I have opportunities to practice skills and procedures more than once.	.306	.392			
CF16 I am having difficulty organizing patient care needs.				.749	
CF5 I am having difficulty prioritizing patient care needs.				.712	
CF17 I feel I may harm a patient due to my lack of knowledge and experience.				.519	
CF8 I feel overwhelmed by my patient care responsibilities and workload.	.226			.457	
CF24 I am experiencing stress in my personal life.				-.210	
CF6 I feel my preceptor provides encouragement and feedback about my work.					.947
CF19 My preceptor is helping me to develop confidence in my practice					.936

Extraction Method: Principal Axis Factoring.

Rotation Method: Promax with Kaiser Normalization.^a

Bolded entries represent items that were retained on each factor.

Figure 4. Model used for confirmatory factor analysis-1 analysis.

Data used for the second CFA-1 testing were the random sample of graduate nurse responses at the six-month time period ($n = 24,378$). The fit statistics used as threshold for acceptance were GFI, NFI, CFI, and RMSEA. Acceptable model fit was defined by the following criteria: GFI (≥ 0.90), NFI (≥ 0.95), RMSEA (≤ 0.06), and CFI (≥ 0.95 ; Hu & Bentler, 1999; Tabachnick & Fidell, 2013). In the first analysis of the model, the Professional Socialization factor (Items CF6 and CF19) was not included as the missing values were problematic for obtaining modification indices. The initial analysis results were chi-square = 30949.544; degrees of freedom = 293; Probability level

= .000. CFI=.858, GFI= .89, RMSEA = .066, NFI= .856. The values for GFI, NFI, and CFI should be above .90 and did not indicate goodness of fit. The standard residual covariance was used to identify pairs of items to co-vary. Items e24 to e28, e21 to e22, and e20 to e21 on the Job Satisfaction factor had high values and were co-varied in the model to provide a better fit. This was labeled Model B. The model fit did improve but did not demonstrate good fit. The standard residual co-variances were examined for items >1.0. Items 26d and 26h on the Job Satisfaction factor were removed. The model was labeled Model C. The fit indices remained the same. It was then decided to remove CF2 (from the support factor) as it had a low loading of .760 on the maximum likelihood estimates. This was labeled Model D and re-run. Then the Professional Socialization factor (two items) was added. This second CFA-1 revealed that overall the indices indicated a good fit to the data, suggesting the five-factor structure, which included a combination of items in Sections II and III, was acceptable. Four items—CF2 (“I am comfortable knowing what to do for a dying patient”), CF26b (“Satisfaction with vacation”), CF26d (“Satisfaction with hours worked”), and CF26h (“Satisfaction with amount of encouragement and feedback”)—were removed, which resulted in a final five-factor structure of 24 items. This CFA-1 confirmed the five-factor structure with 24 items with modifications made was based on theoretical expertise along with using empirical guidelines (see Table 15 for CFA-1 results).

Table 15

Confirmatory Factor Analysis-1 Results Using 24 items and Six-Month Sample

Model	GFI	RMSEA	CFI	NFI	Action Taken
A	.890	.066	.858	.856	Co-vary e24 to e28, e21 to e22, e20 to e21
B	.910	.063	.876	.875	Removed item CF26d
C	.910	.063	.873	.875	Removed items CF26b, CF26h
D	.915	.065	.882	.881	Removed CF2
E		.060	.902	.901	Added Socialization items

Note. $n = 24,378$. GFI = Goodness of fit index, RMSEA = Root mean square error of approximation, CFI = Comparative fit index, NFI = Normed fit index.

A second CFA-2 was conducted to cross-validate the same five-factor structure of 24 items confirmed in the CFA-1 analysis. This confirmatory analysis was conducted on the CFA-2 randomly selected sample of 24,052 graduate nurse responses at the six-month time period. The CFA-2 hypothesized model was drawn in AMOS (see Figure 5 for CFA-2 five-factor structure).

Figure 5. Confirmatory factor analysis-2 final five-factor structure.

The CFA-2 model fit results were as follows: CFI= .897, NFI= .874, and RMSEA = .066. The final five-factor structure with 24 items was confirmed an adequate fit using the CFA-2 data.

Reliability

Reliability estimates for the 24 items in the final five-factor solution were assessed in SPSS. Cronbach alphas for the five subscales—(a) Job Satisfaction, (b) Support, (c) Role Confidence, (d) Ability to Organize/Prioritize Care, and (e)

Professional Socialization—were conducted on the EFA sample ($n = 23,489$), the CFA-1 sample ($n = 24,378$), the CFA-2 sample ($n = 24,052$), and the total sample ($N = 71,919$). Subscale results ranged from .73 to .94, indicating adequate reliability; further analysis showed removing any of the items from the scale did not cause an increase in the alpha value. Results are presented in Table 16 for the identified five factors in the final solution. The total scale reliability estimate using Cronbach's alpha was .86.

Table 16

Reliability Estimates for the Final Five Subscales

Sample	Factor 1-Job Satisfaction	Factor 2 - Support	Factor 3- Role Confidence	Factor 4- Ability to Organize/Prioritize	Factor 5 – Socialization
EFA	.72 ($M = 22.0$)	.83 ($M = 17.33$)	.82 ($M = 21.98$)	.73 ($M = 11.9$)	.94 ($M = 6.84$)
CFA-1	.73 ($M = 22.0$)	.83 ($M = 17.33$)	.82 ($M = 22.0$)	.72 ($M = 11.9$)	.93 ($M = 6.83$)
CFA-2	.73 ($M = 22.0$)	.83 ($M = 17.35$)	.82 ($M = 21.96$)	.73 ($M = 11.9$)	.94 ($M = 6.83$)
Total 6-Month	.73 ($M = 22.0$)	.83 ($M = 17.34$)	.82 ($M = 21.98$)	.73 ($M = 11.9$)	.94 ($M = 6.84$)

Summary

This chapter presented the results of an exploratory factor and confirmatory factor analyses conducted using a large sample collected at the six-month time period from graduate nurses who participated in the Vizient/AACN (Vizient, 2018) 12-month Nurse Residency Program from 2008 to 2018. Characteristics of the sample were presented. The EFA analysis revealed a five-factor solution that was finalized by combining the items in Sections II and II of the survey as those items were based on role transition theory. This five-factor solution explained 49.5% of the total variance in item responses.

The factors were labeled Support, Job Satisfaction, Role Confidence, Organize/Prioritize Care, And Professional Socialization. The five-factor structure was validated by confirmatory analysis and showed an adequate fit of the data to the hypothesized factor structure. Internal consistency was conducted on all responses from each sample. Cronbach alpha for each of the five factor subscales was adequate and ranged from .73 to .94.

CHAPTER V

DISCUSSION AND CONCLUSION

Chapter V presents a discussion of the findings from this study on a factor analysis conducted on the Casey-Fink Survey. A summary of measurement issues encountered during the data analysis, the strengths and limitations of the study, recommendations for future research, and how the study informed revisions to the survey are also reviewed.

Discussion

The purpose of this study was to evaluate the psychometric properties of the Casey-Fink Survey in newly graduated nurses who participated in the Vizient/AACN (Vizient, 2018) 12-month Nurse Residency Program. A secondary data analysis using exploratory factor analysis with a principal axis factoring approach provided strong evidence for a five-factor model, which was confirmed and cross-validated using confirmatory factor analysis methods. This study lays the groundwork to inform future Casey-Fink Survey revisions and refinements and provided additional evidence for the construct validity of the survey.

Previous factor analysis work (Hallaran, 2017; Lynn, 2004; Marshburn, 2007) on the Casey-Fink Survey have shown different factor solutions could be obtained. This study aligned with previous work; however, a different portrait of the factor structure and items retained was drawn. Differences in this factor structure compared to the structure developed by Lynn (2004) might be due to a difference in time period for the assessment

and number of graduate nurses evaluated. This study and the Lynn report used data gathered from graduate nurses who participated in the one-year Vizient/AACN (Vizient, 2018) Nurse Residency Program. Differences in this study's results might reflect the larger sample size, larger number and geographic location of participating healthcare organizations, and a possible difference in the demographics and unit of employment that comprised the sample participants.

Differences in this study's factor structure compared to the Hallaran (2017) study might be due to differences in educational preparation for professional nursing practice, demographic characteristics, use of data at the six-month time period, and type of residency programs offered in Canada.

Major Findings

First, this study confirmed the five-factor solution that emerged from the factor analysis matched the dimensions of the theoretical constructs of role transition. Theory plays a vital role in the development of measurement scales (DeVellis, 2017). Meleis (2010), Kramer (1974), Benner (1984) and Duchscher (2008) all developed theories to understand the process of role transition and the items for the five constructs provided voice for graduate nurses to describe their perceptions of the experience during a role transformation period of time. Benner's novice to expert model clearly supported the importance of a preceptor or mentor to guide the graduate nurse during this advanced beginner stage of entry into practice and skill acquisition. The combination of the 24 items in Section II and nine items from Section III proved to be reflective of the constructs underlying role transition.

Next, only two items on the survey asked about the preceptor's role in encouraging and supporting socialization of the newly graduated nurse into the work unit. These two items loaded strongly on a single factor following factor rotation, which demonstrated they were highly correlated. DeVellis (2017) suggested factors that are represented by two or three items might be underdetermined and might not capture the construct of interest. Empirically, many theories supported the role of a preceptor, especially in the first few months of clinical practice, as a key aspect of successful role transition (Benner, 1984; Duchscher, 2008; Kramer, 1974). Reliability of this subscale was high at .94, reflecting better internal consistency. The two items on this subscale were very important contributors to the content of the scale and were relevant to the construct of role transition, which justified why they were not dropped from the survey. Also problematic was the high number of missing values on these two items at the six-month time period, which was possibly due to respondents not actively working with a preceptor at the time of the survey data collection period. Additional items are needed to better understand this dimension of role transition.

Furthermore, in the past 20 years, the practice environment has become more complex. Measurement of concepts relevant to today's clinical practice environment needs to be included in this survey to ensure its usefulness for healthcare organizations. Content validity is an important aspect of an instrument's item development. Questions related to the use of evidence-based practice decisions need to be included. The knowledge and skills needed for current professional nursing practice has changed significantly since the first version of the Casey-Fink Survey due to a focus on quality outcomes and healthcare cost containment. The Institute of Medicine (2011)

recommended 90% of clinical practice be evidence based. The demand for quality care and patient safety in the current complex healthcare environment requires that nurses use the best possible evidence in their practice to ensure the most beneficial outcomes (Llasus, Angosta, & Clark, 2014). In addition, graduate nurses often work on evidence-based practice projects during their first year of practice.

Five items eliminated during the EFA analysis provided evidence to support survey revisions. Four of the five items loaded on two or more factors and the loading values were less than .4. These questions related to encouragement and feedback provided by the manager, opportunities to practice skills and procedures more than once, and feelings of satisfaction and excitement with work in the chosen nursing specialty.

The stress items were problematic in this analysis for a variety of reasons. Newly graduated nurses encounter multiple work-related stressors that can have serious consequences such as turnover and dissatisfaction with the profession. Understanding what causes stress for graduate nurses is essential to retaining and nurturing this vital workforce entering the nursing profession. Having a single question (CF24) related to concerns about experiences with stress was not sufficient to support the construct of stress in the workplace. This item (CF24) loaded low (-0.210) on Factor 4 (Organize/Prioritize care). This low factor loading in the EFA analysis and the negative correlation did not support keeping this item so it was eliminated. In the follow-up question (CF25), asking respondents to provide a Yes/No (binary choice) response related to the causes of personal stressors. A major shortcoming of binary responses is each item can have only minimal variability (DeVellis, 2017). The fact that the stress items did not load onto any factors in the EFA analysis might have resulted from being a

single question about personal stressors and binary options for the responses in question CF25 led to their removal from the analysis. It has been noted that structured transition to practice programs could improve quality and safety practices, increase job satisfaction, reduce stress, decrease turnover, and also lead to improved patient outcomes (Kramer et al., 2012). Additional items are needed to better measure this aspect of role transition.

During the EFA analysis, four negatively worded items (CF5, CF8, CF16, and CF17) all loaded on the same factor, Organize/Prioritize care. The factor loadings ranged from 0.457 to 0.749. Reversals in item polarity might have been confusing to respondents as they might have failed to notice the difference between expressing their strength of agreement with a statement versus expressing the strength of the attribute being measured (DeVellis, 2017). Future item revisions for this construct need to consider any sources of ambiguity in the wording of the items.

Lastly, of the four items deleted from the final CFA model structure, comfort with caring for a dying patient and amount of feedback and encouragement from the unit manager were single questions on aspects of transition that needed further exploration. The other two items removed were vacation and hours worked, which were related to elements of job satisfaction. Further qualitative studies could provide clarity and understanding of what influenced graduate nurse job satisfaction. These items did not have high factor loadings yet were consistently difficult for graduate nurses during their first year of practice so the items might need additional development and revision.

Strengths and Limitations

Strengths of this study included the very large sample size for conducting psychometric testing ($N = 71,919$). In addition, the graduate nurses included in the

sample closely reflected the diversity in ethnicity and educational backgrounds in today's nursing workforce. In selecting participants for this study, thought was given to whether the research findings could be applied to graduate nurse populations beyond those of the sample that was studied. To increase generalizability of the findings, participants employed in a variety of geographic locations, in different organizational sizes, and participating in various types of nurse residency programs are needed. These research findings were limited to participants from a single nurse residency program curriculum. Furthermore, there might have been differences in BSN versus associate degree nurse responses to the survey.

Convenience samples can suffer from both sampling bias and coverage error. Sampling bias might occur when there are difficulties in getting all the graduate nurses to participate in the study. Coverage error can occur when a portion of the target population is not covered by the sampling frame or the sample does not fit the characteristics of the target population. Therefore, securing new populations of graduate nurses from a variety of practice settings and geographic areas is needed to increase the generalizability of study findings and to obtain perceptions from non-proprietary nurse residency programs and from graduate nurses working in a variety of clinical practice environments.

Another limitation of this study involved the use of a self-report instrument, which could have led to bias. A potential threat to internal validity includes selection bias as the sample in this study only included newly graduated nurses who participated in the Vizient/AACN (Vizient, 2018) 12-month Nurse Residency Program. Moreover, failure of participants to respond to all the questions on the survey instrument, especially

with long questionnaires, could also threaten the validity of the instrument (Gray et al., 2017).

Recommendations for Research

As with all survey instruments, the Casey-Fink Survey needs to be refined and updated to include new knowledge and competencies required for graduate nurses to practice safely. Items that address perceptions of current practice environment issues, confidence implementing evidence in practice, identifying workplace stressors that impact turnover, and other potential aspects of the transition experience that might not be represented in the current survey items need to be developed. More research is needed on what factors affect graduate nurses' job satisfaction as they might differ from those of experienced nurses. Additional research is needed on scale development items. This would require direct input about item creation from potential respondents. DeVellis (2017) suggested qualitative research methods to study facts, observations, and experiences could be used as empirical indicators when developing an instrument (Knapp, 1998). Focus groups of current graduate nurses might be useful for identifying empirical indicators of role transition. The use of cognitive interviewing techniques to better understand new items and gather input and perceptions about the role transition experience from current graduate nurses during their residency program are other appropriate and useful strategies in new item development. Additional validation studies are needed using different samples of graduate nurses working in internally developed nurse residency programs in multiple hospital sites. Selection of a different time period during the first year of practice might provide different or similar results. Because the

factor structure was easily influenced by sampled data, repetitive revalidation studies are needed to overcome sampling bias and to confirm the latent variable structure.

This study revealed this survey did not adequately measure perceived levels of job-related stress and stressors in the clinical environment. To address this problem, future research needs to identify graduate nurse perceived stress and stressors in the clinical practice environment so they can be accurately measured. Newly graduated nurses could be targets of incivility because of their lack of confidence and experience. Understanding the population for whom the scale is intended is critical for developing good items (Polit & Beck, 2012). Themes identified from a qualitative study assessing the use of mentors to guide professional growth beyond the preceptor relationship and to identify perceived stress and stressors in the workplace could be utilized to develop additional items on what is stressful for graduate nurses during their role transition. These items would need to be pilot tested to provide a larger number of questions to be included in future survey revisions.

Future research that includes examining the factor structure of the Casey-Fink Survey at the baseline and 12-month data collection time periods is needed. Testing the differences between graduate nurse demographic characteristics and the five constructs identified in this study at all three data collection time periods using the database of responses provided by Vizient/AACN (Vizient, 2018) would be another area of study.

Conclusion

This study described the steps in testing the factor structure of Casey-Fink Survey using a large sample of graduate nurses. In this study, graduate nurse role transition constructs were inferred and validated by conducting exploratory and confirmatory factor

analysis on data collected from items in Sections II and III of the Casey-Fink Survey. This study added new information to the literature about psychometric testing of the Casey-Fink Survey instrument. The Casey-Fink Survey has remained in use throughout the Vizient/AACN (Vizient, 2018) program's history and continues to be valuable in the evaluation and design of nurse residency programs. Revisions to the survey must reflect current clinical practice environments. A good-fitting model for the Casey-Fink Survey could still be improved upon and tested to link role transition and retention outcomes. Examination of the phenomenon of graduate nurse transition into practice is a significant area of study in nursing because it affects nurse retention. It is essential that measures of graduate nurse role transition are both current and psychometrically sound. Additional validation studies are needed that (a) use a variety of populations of graduate nurses who work in diverse healthcare organization settings and (b) measure graduate nurse role transition with the Casey-Fink Survey in an assortment of transition programs. Accurate data are needed to design and evaluate graduate nurse transition into practice and because the Casey-Fink Survey has been used to collect this data, it was vital to determine the psychometric properties of the tool. A validation study on the reliability and structural validity of the concepts measured by the Casey-Fink Survey was a significant contribution to the body of nursing's knowledge on measurement instruments specific to graduate nurse role transition. In addition, this study was needed to inform revisions to the Casey-Fink Survey.

REFERENCES

- American Nurses Credentialing Center. (2018). *Practice transition accreditation program manual*. Silver Spring, MD: American Nurses Credentialing Center.
- Anderson, G., Hair, C., & Toderro, C. (2012). Nurse residency programs: An evidence-based review of theory, process, and outcomes. *Journal of Professional Nursing*, 28(4), 203-212.
- Benner, P. (1984). *From novice to expert: Excellence and power in clinical nursing practice*. Menlo Park, CA: Addison-Wesley Publishing.
- Bowles, C., & Candela, L. (2005). First job experiences of recent RN graduates: Improving the work environment. *Journal of Nursing Administration*, 35, 130-137.
- Brewer, C. S., Kovner, C. T., Greene, W., Tukov-Shuser, M., & Djukic, M. (2012). Predictors of actual turnover in a national sample of newly licensed registered nurses employed in hospitals. *Journal of Advanced Nursing*, 68(3), 521-538.
- Brown, T. A. (2015). *Confirmatory factor analysis for applied research* (2nd ed.). New York, NY: Guilford Press.
- Casey, K., Fink, R., Krugman, M., & Propst, J. (2004). The graduate nurse experience. *Journal of Nursing Administration*, 34(6), 303-311.

- Cline, D., La Frentz, K., Fellman, B., Summers, B., & Brassil, K. (2017). Longitudinal outcomes of an institutionally developed nurse residency program. *Journal of Nursing Administration*, 47(7/7), 384-390.
doi:10.1097/NNA.0000000000000500
- Collaborative Institutional Training Initiative. (2017). *Research ethics and compliance training*. Retrieved from <https://www.citiprogram.org>
- Commission on Collegiate Nursing Education. (2015). *Standards for accreditation of entry-to-practice nurse residency programs*. Retrieved from <http://www.aacn.nche.edu/ccne-accreditation/CCNE-Entry-to-Practice-Residency-Standards-2015.pdf>.
- Costello, A. B., & Osborne, J. W. (2005). Best practices in exploratory factor analysis: Four recommendations for getting the most from your analysis. *Practical Assessment, Research & Evaluation*, 10(7). Retrieved from <http://pareonline.net/getvn.asp?v=10&n=7>
- Cylke, K. (2012). Needs assessment. *Journal for Nurses in Staff Development*, 28(2), 57-61.
- DeVellis, R. F. (2017). *Scale development: Theory and applications* (4th ed.). Thousand Oaks: SAGE.
- Dubois, C. A., & Zedreck Gonzales, J. F. (2018). Implementing a resilience-promoting education program for new nursing graduates. *Journal for Nurses in Professional Development*, 34(5), 263-269.

- Duchscher, J. E. B. (2008). A process of becoming: The stages of new nursing graduate professional role transition. *The Journal of Continuing Education in Nursing*, 39(10), 441-450.
- Duchscher, J. B., & Windey, M. (2018). Stages of transition and transition shock. *Journal for Nurses in Professional Development*, 34(4), 228-232.
doi:10.1097/NND.0000000000000461
- Fink, R., Krugman, M., Casey, K., & Goode, C. (2008). The graduate nurse experience: Qualitative residency program outcomes. *Journal of Nursing Administration*, 38(7/8), 341-348.
- Friday, L., Zoller, J. S., Hollerbach, A. D., Jones, K., & Knofczynski, G. (2015). The effects of a pre-licensure extern program and nurse residency program on new graduate outcomes and retention. *Journal for Nurses in Professional Development*, 31(3), 151-157. doi:10.1097/NND.0000000000000158
- Goode, C. J., Lynn, M. L., Krsek, C., & Bednash, G. D. (2009). Nurse residency programs: An essential requirement for nursing. *Journal of Nursing Administration*, 27(3), 142-147.
- Goode, C. J., Lynn, M. R., McElroy, D., Bednash, G. D., & Murray, B. (2013). Lessons learned from 10 years of research on a post-baccalaureate nurse residency program. *Journal of Nursing Administration*, 43(2), 71-77.
- Goode, C. J., Ponte, P. R., & Havens, D. S. (2016). Residency for transition into practice. *The Journal of Nursing Administration*, 46(2), 82-86.

- Gray, J. R., Grove, S. K., & Sutherland, S. (2017). *The practice of nursing research: Appraisal, synthesis, and generation of evidence* (8th ed.). St. Louis, MO: Elsevier.
- Grove, S. K., Gray, J. R., & Burns, N. (2015). *Understanding nursing research: Building an evidence-based practice* (6th ed.). St. Louis, MO: Elsevier-Saunders.
- Hallaran, A. J. (2017). *Understanding the variables influencing graduate nurse transition and retention in the workplace*. Retrieved from https://qspace.library.queensu.ca/bitstream/handle/1974/15913/Hallaran_Amy_J_201706_PHD.pdf?sequence=2&isAllowed=y
- Harrison, D., & Ledbetter, C. (2014). Nurse residency programs. *Journal for Nurses in Professional Development*, 30(2), 76-82.
- Hillman, L., & Foster, R. R. (2011). The impact of a nursing transitions programme on retention and cost savings. *Journal of Nursing Management*, 19(1). 50-56.
doi:10.1111/j.1365-2834.2010.01187.x
- Hu, L. T., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal*, 6(1), 1-55.
- Institute of Medicine. (2011). *The future of nursing: Leading change, advancing health*. Washington, DC: National Academies Press.
- Johnson, A., Salisbury, H., Johannsson, M., & Barajas, K. (2013). Emergency nurse residency program evaluation, *Journal for Nurses in Professional Development*, 29(5), 233-237.
- Kaiser, H. F. (1974). An index of factorial simplicity. *Psychometrika*, 39(1), 31-36.

- Kline, R. B. (2011). *Principles and practice of structural equation modeling*. New York, NY: Guilford Press.
- Klingbeil, C., Schiffman, R. F., Ziebert, C., Totka, J. P., Schmitt, C. A., Doyle, L., ... Johnson, N. (2016). Transition of experienced and new graduate nurses to a pediatric hospital. *Journal for Nurses in Professional Development*, 32(4), 198-204.
- Knapp, T. R. (1998). *Quantitative nursing research*. Thousand Oaks, CA: Sage.
- Kovner, C. T., Brewer, C. S., Fatehi, F., & Jun, J. (2014). What does nurse turnover rate mean and what is the rate? *Policy, Politics & Nursing Practice*, 15(3-4), 64-71.
doi:10.1177/1527154414547953
- Kovner, C. T., & Djukic, M. (2009). The nursing career process from application through the first two years of employment. *Journal of Professional Nursing*, 25(4), 197-203.
- Kramer, M. (1974). *Reality shock: Why nurses leave nursing*. St. Louis, MO: C.V Mosby.
- Kramer, M., Maguire, P., Schmalenberg, C., Halfer, D. S., Budin, W. C., Hall, D. S., ... Leme, J. (2012). Components and strategies of nurse residency programs effective in new graduate socialization. *Western Journal of Nursing Research*, 35(5), 566-589.
- Krugman, M., Bretschneider, J., Horn, P. B., Kresk, C. A., Moutafis, R. A., & Smith, M. O. (2006). The national post-baccalaureate graduate nurse residency program: A model for excellence in transition to practice. *Journal for Nurses in Staff Development*, 22(4), 196-204.

- Llasus, L., Angosta, A. D., & Clark, M. (2014). Graduating baccalaureate students' evidence-based practice knowledge, readiness, and implementation. *Journal of Nursing Education, 53*(9), S82- S89.
- Letourneau, R. M., & Fater, K. H. (2015). Nurse residency programs: An integrative review of the literature. *Nursing Education Perspectives, 36*(2), 96-101.
- Lynn, M. R. (2004). *Summary of UHC analysis of Casey-Fink graduate nurse experience survey*. Prepared for Regina Fink March 22, 2004.
- Marshburn, D. M. (2007). *Clinical competence, satisfaction, and intent to stay in new nurses* (Doctoral dissertation). East Carolina University, Greenville, North Carolina.
- Marshburn, D. M., Engelke, M. K., & Swanson, M. S. (2009). Relationships of new nurses' perceptions and measured performance-based clinical competence. *The Journal of Continuing Education in Nursing, 40*(9), 426-432.
- Medas, J. C., Amato, S., Grimm, D., Radziewicz, R., Rhodes, C., VanHorn, C., & McNett, M. (2015). Outcomes of a comprehensive nurse residency program. *Nursing Management, 46*(11), 41-48.
- Meleis, A. I. (2010). *Transitions theory: Middle-range and situation-specific theories in nursing research and practice*. New York, NY: Springer Publishing Company.
- Meleis, A. I., Sawyer, L., Im, E. O., Hilfinger, M. D., & Schumacher, K. (2000). Experiencing transitions: An emerging middle-range theory. *Advances in Nursing Science, 23*(1), 12-28.
- Meyer, G., & Shatto, B. (2017). Resilience and transition to practice in direct entry nursing graduates. *Nurse Education in Practice, 28*, 276-279.

- Nunnally, J. C. (1978). *Psychometric theory* (2nd ed.). New York, NY: McGraw-Hill.
- Olson-Sitki, K., Wendler, M. C., & Forbes, G. (2012). Evaluating the impact of a nurse residency program for newly graduated registered nurses. *Journal for Nurses in Staff Development*, 28(4), 156-162. doi:10.1097/NND.0b013e31825dfb4c
- Pett, M. A., Lackey, N. R., & Sullivan, J. J. (2003). *Making sense of factor analysis: The use of factor analysis for instrument development in health care research*. Thousand Oaks, CA: Sage Publications.
- Polit, D. F. (2010). *Statistics and data analysis for nursing research* (2nd ed.). Upper Saddle River, NJ: Pearson Education.
- Polit, D. F., & Beck, C. T. (2012). *Nursing research: Generating and assessing evidence for nursing practice* (9th ed.). Philadelphia, PA: Lippincott Williams, & Wilkins.
- Polit, D. F., & Yang, F. M. (2016). *Measurement and the measurement of change*. Philadelphia: Wolters Kluwer.
- Rush, K. L., Adamack, M., Janke, R., Gordon, J., & Ghement, I. R. (2012). The helpfulness and timing of transition program education. *Journal for Nurses in Professional Development*, 29(4), 191-196.
- Schumacher, K., & Meleis, A. I. (1994). Transitions: A central concept in nursing. *Journal of Nursing Scholarship*, 26(2), 119-127.
- Slate, K. A., Stavarski, D. H., Romig, B. J., & Thacker, K. S. (2018). Longitudinal study transformed onboarding nurse graduates. *Journal for Nurses in Professional Development*, 34(2), 92-98.

- Soeken, K. L. (2010). Validity of measures. In C. F. Waltz, O. L. Strickland, & E. R. Lenz (Eds.), *Measurement in nursing and health research* (pp. 163-201). New York: Springer.
- Stephenson, J. K., & Cosme, S. (2018). Instruments to evaluate nurse residency programs. *Journal for Nurses in Professional Development*, 34(3), 123-132.
- Tabachnick, B. G., & Fidell, L. S. (2013). *Using multivariate statistics* (6th ed.). Boston: Pearson Education.
- Thomson, S. (2011). Transition into practice: A comparison of outcomes between associate- and baccalaureate-prepared nurses participating in a nurse residency program. *Journal for Nurses in Staff Development*, 27(6), 266-271.
- Tran, U. S., & Formann, A. K. (2009). Performance of parallel analysis in retrieving unidimensionality in the presence of binary data. *Educational and Psychological Measurement*, 69, 5-61.
- Tyo, M. B., Gundlach, M., Brennan, C., Esdale, L., Knight, A., Provencher, S., & Tardy, K. (2018). Leading the charge: Achievement of national accreditation for a nurse residency program. *Journal for Nurses in Professional Development*, 34(5), 270-276.
- Vizient. (2018). *AACN nurse residency program: Our solutions*. Retrieved from <https://www.vizientinc.com/Our-solutions/Clinical-Solutions/Vizient-AACN-Nurse-Residency-Program>.
- Wenger, B. (2015). Development of an intravenous skills module for graduate nurses. *Infusion Nurses Society*, 38(2), 135-139.

- Williams, B., Brown, T., & Onsman, A. (2010). Exploratory factor analysis: A five-step guide for novices. *Australasian Journal of Paramedicine*, 8(3), 1-13.
- Wilson, T., Weathers, N., & Forneris, L. (2018). Evaluation of outcomes from an online nurse residency program. *The Journal of Nursing Administration*, 48(10), 495-501.

APPENDIX A
INSTITUTIONAL REVIEW BOARD APPROVAL

Institutional Review Board

DATE: June 27, 2019

TO: Kathryn Casey, PhD

FROM: University of Northern Colorado (UNCO) IRB

PROJECT TITLE: [1434846-1] Psychometric Evaluation of the Casey-Fink Graduate Nurse Experience Survey

SUBMISSION TYPE: New Project

ACTION: APPROVAL/VERIFICATION OF EXEMPT STATUS

DECISION DATE: June 27, 2019

EXPIRATION DATE: June 27, 2023

Thank you for your submission of New Project materials for this project. The University of Northern Colorado (UNCO) IRB approves this project and verifies its status as EXEMPT according to federal IRB regulations.

We will retain a copy of this correspondence within our records for a duration of 4 years.

If you have any questions, please contact Nicole Morse at 970-351-1910 or nicole.morse@unco.edu. Please include your project title and reference number in all correspondence with this committee.

This letter has been electronically signed in accordance with all applicable regulations, and a copy is retained within University of Northern Colorado (UNCO) IRB's records.

APPENDIX B
NON-DISCLOSURE AND CONFIDENTIALITY
AGREEMENT

**VIZIENT
NON-DISCLOSURE AND
CONFIDENTIALITY AGREEMENT**

THIS Non-Disclosure and Confidentiality Agreement (“**Agreement**”) is between Vizient, Inc., on behalf of itself and its subsidiaries (collectively, “**Vizient**”), a Delaware limited liability company with offices at 155 North Wacker Drive, Chicago, IL 60606, and **Kathryn Kathryn C. Casey**, located at **2548 Ward Drive, Lakewood CO 80215** (“**Individual**”). Vizient and Individual are singularly referred to as “**Party**” and collectively referred to as “**Parties**”. This Agreement is dated **November 23, 2018**, 2018 (“**Effective Date**”).

RECITALS

WHEREAS, Vizient and Individual each take the obligation to protect Vizient’s confidential, proprietary and commercially valuable information very seriously, and enter into this Agreement with the specific intent of protecting the Confidential Information, as defined below;

WHEREAS, Vizient may disclose to Individual certain confidential information in connection with research collaboration described herein (“**Research Collaboration**”) and each desires that any such information shall be kept confidential by Individual;

WHEREAS, in consideration of Vizient’s disclosure, Individual is willing to hold in trust Vizient’s Confidential Information and keep it strictly confidential in accordance with the terms and conditions set forth in this Agreement; and

NOW, THEREFORE, for the mutual premises contained herein and other valuable consideration, the sufficiency of which is hereby acknowledged, Vizient and Individual hereby agree as follows:

1. DEFINITIONS:

Confidential Information shall mean and include, but is not limited to: the information of Vizient, its member organizations (“**Members**”), and other participants enrolled in the Nurse Residency Program (“**NRP**”) and each Member’s and participant’s NRP data and all other related data bases, studies, project data, technology, plans, designs, concepts, computer programs, videos, animation and designs; computer codes, including but not limited to, HTML, .NET, ASP, SQL, JAVA or Javascript; formulas and websites.

“**Research Collaboration**” means Individual’s access to confidential information/de-identified information for purposes of using Vizient Casey-Fink survey results for Individual’s PhD dissertation project, including conducting a factor analysis for possible revisions to the survey instrument.

2. RESTRICTIONS AND OBLIGATIONS:

a. Confidential Information furnished, directly or indirectly, in writing or otherwise, to Individual by Vizient under this Agreement is and shall remain the property of Vizient, and shall be used only as provided herein.

b. Individual agrees to hold in confidence any Confidential Information of Vizient to which it has access and shall not utilize, except for the Research Collaboration provided herein, or disclose such Confidential Information to any person or entity. Confidential Information includes any materials produced in writing which are clearly identified as confidential, and any oral disclosures where Vizient has indicated at the time of disclosure the confidential or proprietary nature of the information or which the Individual should reasonably know to be the confidential information of Vizient.

c. Confidential Information shall not include: (a) information known to Individual prior to receiving such information from Vizient; (b) information rightfully obtained from a third party without similar

restrictions and without breach of this Agreement; (c) information independently developed by Individual; or (d) information publicly available.

d. Vizient expressly denies Individual permission to otherwise use, disclose or in any other manner copy any Confidential Information Individual may have seen or been shown at any time, at or by Vizient for the Business Purpose, without the written permission of Vizient;

e. Individual agrees to share a copy of their dissertation paper upon completion with Vizient, or any other paper intended for publication that included Vizient Confidential Information;

f. This section shall survive termination of this Agreement.

3. REMEDIES:

The Parties agree that, in the event of breach of this Agreement by Individual, it may be difficult for Vizient to establish the amount of loss or damages with reasonable certainty. The Individual agrees that, in addition to other legal relief that may be available to Vizient, this Agreement may be sought to be enforced by injunctive relief.

4. TERM and TERMINATION

The Parties agree that the term of this Agreement shall expire at project end. All rights and obligations of this agreement shall survive expiration for a period of three (3) years from the Effective Date.

IN WITNESS WHEREOF, Individual and Vizient have each caused this Agreement to be signed all as of the Effective Date.

SO AGREED.

VIZIENT, INC., ON BEHALF OF ITSELF AND ITS SUBSIDIARIES

By: Jayne Willingham

Printed Name: _____

Title: _____

Date: _____

INDIVIDUAL

By: Kathryn Casey

Printed Name: Kathryn Casey

Title: PhD Student University of Northern Colorado

Date: November 23, 2018

APPENDIX C
CASEY-FINK GRADUATE NURSE
EXPERIENCE SURVEY

Casey-Fink Graduate Nurse Experience Survey (revised)

© 2006 University of Colorado Hospital. All rights reserved.

-
- I. List the top three skills/procedures you are *uncomfortable performing* independently at this time? (please select from the drop down list) **list is at the end of this document.****

1. _____
 2. _____
 3. _____
 4. _____ I am independent in all skills
-

II. Please answer each of the following questions by placing a mark inside the circles:

	STRONGLY DISAGREE	DISAGREE	AGREE	STRONGLY AGREE
1. I feel confident communicating with physicians.	☐	☐	☐	☐
2. I am comfortable knowing what to do for a dying patient.	☐	☐	☐	☐
3. I feel comfortable delegating tasks to the Nursing Assistant.	☐	☐	☐	☐
4. I feel at ease asking for help from other RNs on the unit.	☐	☐	☐	☐
5. I am having difficulty prioritizing patient care needs.	☐	☐	☐	☐
6. I feel my preceptor provides encouragement and feedback about my work.	☐	☐	☐	☐
7. I feel staff is available to me during new situations and procedures.	☐	☐	☐	☐
8. I feel overwhelmed by my patient care responsibilities and workload.	☐	☐	☐	☐
9. I feel supported by the nurses on my unit.	☐	☐	☐	☐
10. I have opportunities to practice skills and procedures more than once.	☐	☐	☐	☐
11. I feel comfortable communicating with patients and their families.	☐	☐	☐	☐

	STRONGLY DISAGREE	DISAGREE	AGREE	STRONGLY AGREE
12. I am able to complete my patient care assignment on time.	☐	☐	☐	☐
13. I feel the expectations of me in this job are realistic.	☐	☐	☐	☐
14. I feel prepared to complete my job responsibilities.	☐	☐	☐	☐
15. I feel comfortable making suggestions for changes to the nursing plan of care.	☐	☐	☐	☐
16. I am having difficulty organizing patient care needs.	☐	☐	☐	☐
17. I feel I may harm a patient due to my lack of knowledge and experience.	☐	☐	☐	☐
18. There are positive role models for me to observe on my unit.	☐	☐	☐	☐
19. My preceptor is helping me to develop confidence in my practice.	☐	☐	☐	☐
20. I am supported by my family/friends.	☐	☐	☐	☐
21. I am satisfied with my chosen nursing specialty.	☐	☐	☐	☐
22. I feel my work is exciting and challenging.	☐	☐	☐	☐
23. I feel my manager provides encouragement and feedback about my work.	☐	☐	☐	☐
24. I am experiencing stress in my personal life.	☐	☐	☐	☐

25. If you chose agree or strongly agree, to #24, please indicate what is causing your stress. (You may circle more than once choice.)

- a. NCLEX
- b. Finances
- c. Child care
- d. Living situation
- e. Personal relationships
- f. Job performance
- g. Graduate school

III. How *satisfied* are you with the following aspects of your job:

	VERY DISSATISFIED	MODERATELY DISSATISFIED	NEITHER SATISFIED NOR DISSATISFIED	MODERATELY SATISFIED	VERY SATISFIED
Salary	☐	☐	☐	☐	☐
Vacation	☐	☐	☐	☐	☐
Benefits package	☐	☐	☐	☐	☐
Hours that you work	☐	☐	☐	☐	☐
Weekends off per month	☐	☐	☐	☐	☐
Your amount of responsibility	☐	☐	☐	☐	☐
Opportunities for career advancement	☐	☐	☐	☐	☐
Amount of encouragement and feedback	☐	☐	☐	☐	☐
Opportunity to work straight days	☐	☐	☐	☐	☐

IV. Transition (please circle any or all that apply)**1. What difficulties, if any, are you currently experiencing with the transition from the "*student*" role to the "*RN*" role?**

- a. role expectations (e.g. autonomy, more responsibility, being a preceptor or in charge)
- b. lack of confidence (e.g. MD/PT communication skills, delegation, knowledge deficit, critical thinking)
- c. workload (e.g. organizing, prioritizing, feeling overwhelmed, ratios, patient acuity)
- d. fears (e.g. patient safety)
- e. orientation issues (e.g. unit familiarization, learning technology, relationship with multiple preceptors, information overload)

2. What could be done to help you feel more supported or integrated into the unit?

- a. improved orientation (e.g. preceptor support and consistency, orientation extension, unit specific skills practice)
- b. increased support (e.g. manager, RN, and educator feedback and support, mentorship)
- c. unit socialization (e.g. being introduced to staff and MDs, opportunities for staff socialization)
- d. improved work environment (e.g. gradual ratio changes, more assistance from unlicensed personnel, involvement in schedule and committee work)

3. What aspects of your work environment are most satisfying?

- a. peer support (e.g. belonging, team approach, helpful and friendly staff)
- b. patients and families (e.g. making a difference, positive feedback, patient satisfaction, patient interaction)
- c. ongoing learning (e.g. preceptors, unit role models, mentorship)
- d. professional nursing role (e.g. challenge, benefits, fast pace, critical thinking, empowerment)
- e. positive work environment (e.g. good ratios, available resources, great facility, up-to-date technology)

4. What aspects of your work environment are least satisfying?

- a. nursing work environment (e.g. unrealistic ratios, tough schedule, futility of care)
- b. system (e.g. outdated facilities and equipment, small workspace, charting, paperwork)
- c. interpersonal relationships (e.g. gossip, lack of recognition, lack of teamwork, politics)
- d. orientation (inconsistent preceptors, lack of feedback)

5. Please share any comments or concerns you have about your residency program:

Demographics: Circle the response that represents the most accurate description of your individual professional profile.

1. Age: _____ years

2. Gender:

- a. Female
- b. Male

3. Ethnicity:

- a. Caucasian (white)
- b. Black
- c. Hispanic
- d. Asian
- e. Other
- f. I do not wish to include this information

4. Area of specialty:

- a. Adult Medical/Surgical
- b. Adult Critical Care
- c. OB/Post-Partum
- d. NICU
- e. Pediatrics
- f. Emergency Department
- g. Oncology
- h. Transplant
- i. Rehabilitation
- j. OR/PACU
- k. Psychiatry
- l. Ambulatory Clinic
- m. Other: _____

5. School of Nursing Attended (name, city, state located): _____**6. Date of Graduation:** _____**7. Degree Received:** AD: _____ Diploma: _____ BSN: _____ ND: _____**8. Other Non-Nursing Degree (if applicable):** _____**9. Date of Hire (as a Graduate Nurse):** _____**10. What previous health care work experience have you had:**

- a. Volunteer
- b. Nursing Assistant
- c. Medical Assistant
- d. Unit Secretary
- e. EMT
- f. Student Externship
- g. Other (*please specify*): _____

11. Have you functioned as a charge nurse?

- a. Yes
- b. No

12. Have you functioned as a preceptor?

- a. Yes
- b. No

13. What is your scheduled work pattern?

- a. Straight days
- b. Straight evenings
- c. Straight nights
- d. Rotating days/evenings
- e. Rotating days/nights
- f. Other (*please specify*): _____

14. How long was your unit orientation?

- a. Still ongoing
- b. ≤ 8 weeks
- c. 9 – 12 weeks
- d. 13 – 16 weeks
- e. 17 - 23 weeks
- f. ≥ 24 weeks

15. How many *primary* preceptors have you had during your orientation?

_____ **Number of preceptors**

16. Today's date: _____**Drop down list of skills**

Arterial/venous lines/swan ganz (wedging, management, calibration, CVP, cardiac output)

Assessment skills

Bladder catheter insertion/irrigation

Blood draw/venipuncture

Blood product administration/transfusion

Central line care (dressing change, blood draws, discontinuing)

Charting/documentation

Chest tube care (placement, pleurovac)

Code/Emergency Response

Death/Dying/End-of-Life Care

Dobhoff/NG care/suctioning/placement

ECG/EKG/Telemetry monitoring and interpretation

Intravenous (IV) medication administration/pumps/PCAs

Intravenous (IV) starts

Medication administration

MD communication

Patient/family communication and teaching

Prioritization/Time Management

Trach care

Vent care/management/assisting with intubation/extubation

Wound care/dressing change/wound vac

Unit specific skills _____